

**B4GALT1 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8892b**

### Specification

#### B4GALT1 Antibody (C-term) - Product Information

Application	WB, FC,E
Primary Accession	<a href="#">P15291</a>
Other Accession	<a href="#">P15535</a> , <a href="#">P08037</a>
Reactivity	Human
Predicted	Bovine, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	329-358

#### B4GALT1 Antibody (C-term) - Additional Information

##### Gene ID 2683

##### Other Names

Beta-1, 4-galactosyltransferase 1, Beta-1, 4-GalTase 1, Beta4Gal-T1, b4Gal-T1, 241-, UDP-Gal:beta-GlcNAc beta-1, 4-galactosyltransferase 1, UDP-galactose:beta-N-acetylglucosamine beta-1, 4-galactosyltransferase 1, Lactose synthase A protein, N-acetyllactosamine synthase, Nal synthase, Beta-N-acetylglucosaminylglycopeptide beta-1, 4-galactosyltransferase, Beta-N-acetylglucosaminyl-glycolipid beta-1, 4-galactosyltransferase, 241-, Processed beta-1, 4-galactosyltransferase 1, B4GALT1, GGTB2

##### Target/Specificity

This B4GALT1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 329-358 amino acids from the C-terminal region of human B4GALT1.

##### Dilution

WB~~1:1000

FC~~1:10~50

E~~Use at an assay dependent concentration.

##### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

##### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions

B4GALT1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## B4GalT1 Antibody (C-term) - Protein Information

**Name** B4GALT1 ([HGNC:924](#))

**Synonyms** GGTB2

**Function** Galactosyltransferase acting in the Golgi stacks. Catalyzes the transfer of galactose (Gal) from UDP-alpha-D-galactose in beta(1->4) linkage to the non-reducing terminal N-acetylglucosamine (GlcNAc) moieties of glycolipids and complex-type N-linked glycans (PubMed:[16157350](#), PubMed:[27872474](#), PubMed:[29133956](#), PubMed:[36280670](#), PubMed:[37632720](#), PubMed:[38321209](#)). Adds one Gal residue to both GlcNAc beta(1->2)-linked to the alpha(1->3) and alpha(1->6) mannose antennae of complex-type N-glycans, enabling the formation of mono- and di- galactosylated glycoforms. Galactosylates complex-type N-glycans attached on the fragment crystallizable (Fc) of immunoglobulin-gamma isotypes (IgGs), a prerequisite for antibody glycan sialylation and related anti-inflammatory effector functions (PubMed:[27872474](#), PubMed:[29133956](#), PubMed:[36280670](#), PubMed:[37632720](#)). Can also transfer a Gal residue to free GlcNAc to form N-acetyllactosamine (PubMed:[16157350](#)). With LALBA/alpha-lactalbumin forms the lactose synthase complex responsible for production of large amounts of lactose in the lactating mammary gland. Interaction with LALBA alters the sugar substrate specificity of the catalytic domain, enabling high affinity binding of glucose and its transformation to lactose (PubMed:[16157350](#)).

### Cellular Location

Golgi apparatus, Golgi stack membrane; Single-pass type II membrane protein Note=Colocalizes with MGAT3 and ST6GAL1 in the Golgi stacks of cisternae, including the cis-part of the Golgi stacks [Isoform Short]: Golgi apparatus, Golgi stack membrane; Single-pass type II membrane protein. Note=Found in trans cisternae of Golgi

### Tissue Location


Ubiquitously expressed, but at very low levels in fetal and adult brain


## B4GalT1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## B4GalT1 Antibody (C-term) - Images





All lanes : Anti-B4GalT1 Antibody (C-term) at 1:1000 dilution Lane 1: HeLa, whole cell lysate Lane 2: T47D whole cell lysate ysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 50 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



B4GalT1 Antibody (C-term) (Cat. #AP8892b) flow cytometric analysis of HL-60 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### **B4GalT1 Antibody (C-term) - Background**

B4GalT1 is an enzyme that participates both in glycoconjugate and lactose biosynthesis. For the first activity, the enzyme adds galactose to N-acetylglucosamine residues that are either monosaccharides or the nonreducing ends of glycoprotein carbohydrate chains. The second activity is restricted to lactating mammary tissues where the enzyme forms a heterodimer with alpha-lactalbumin to catalyze UDP-galactose + D-glucose UDP + lactose. The two enzymatic forms result from alternate transcription initiation sites and post-translational processing.

#### **B4GalT1 Antibody (C-term) - References**

Mengle-Gaw,L., et.al., Biochem. Biophys. Res. Commun. 176 (3), 1269-1276 (1991)