

**PMEL17 / GP100 Antibody**  
**Rabbit mAb**  
**Catalog # AP90205****Specification**

---

**PMEL17 / GP100 Antibody - Product Information**

|                                                                                  |                        |
|----------------------------------------------------------------------------------|------------------------|
| Application                                                                      | WB, IHC                |
| Primary Accession                                                                | <a href="#">P40967</a> |
| Clonality                                                                        | Monoclonal             |
| <b>Other Names</b>                                                               |                        |
| PMEL; ME20-M; Premelanosome protein; Silver locus protein homolog; PMEL17; SILV; |                        |
| Isotype                                                                          | Rabbit IgG             |
| Host                                                                             | Rabbit                 |
| Calculated MW                                                                    | 70255 Da               |

**PMEL17 / GP100 Antibody - Additional Information**

|                              |                                                                                                                                                                                                                                                                                                                                                                                         |
|------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Dilution                     | WB~~1:1000<br>IHC~~1:100~500                                                                                                                                                                                                                                                                                                                                                            |
| Purification                 | Affinity-chromatography                                                                                                                                                                                                                                                                                                                                                                 |
| Immunogen                    | A synthesized peptide derived from human PMEL17 / GP100                                                                                                                                                                                                                                                                                                                                 |
| Description                  | Plays a central role in the biogenesis of melanosomes. Involved in the maturation of melanosomes from stage I to II. The transition from stage I melanosomes to stage II melanosomes involves an elongation of the vesicle, and the appearance within of distinct fibrillar structures. Release of the soluble form, ME20-S, could protect tumor cells from antibody mediated immunity. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.                                                                                                                                                                                                       |

**PMEL17 / GP100 Antibody - Protein Information****Name** PMEL**Synonyms** D12S53E, PMEL17, SILV**Function**

Forms physiological amyloids that play a central role in melanosome morphogenesis and pigmentation. The maturation of unpigmented premelanosomes from stage I to II is marked by assembly of processed amyloidogenic fragments into parallel fibrillar sheets, which elongate the

vesicle into a striated ellipsoidal shape. In pigmented stage III and IV melanosomes, the amyloid matrix serves as a platform where eumelanin precursors accumulate at high local concentrations for pigment formation. May prevent pigmentation-associated toxicity by sequestering toxic reaction intermediates of eumelanin biosynthesis pathway.

#### **Cellular Location**

Endoplasmic reticulum membrane; Single-pass type I membrane protein. Golgi apparatus, cis-Golgi network membrane; Single-pass type I membrane protein. Endosome, multivesicular body. Melanosome Extracellular vesicle. Secreted. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065) Localizes predominantly to intraluminal vesicles (ILVs) within multivesicular bodies. Associates with ILVs found within the lumen of premelanosomes and melanosomes and particularly in compartments that serve as precursors to the striated stage II premelanosomes (PubMed:11694580, PubMed:12643545). Sorted to stage I melanosomes following its processing in the ER and cis-Golgi (PubMed:15096515) Transiently expressed at the cell surface before targeting to early melanosomes (PubMed:16760433, PubMed:30988362). Colocalizes with BACE2 in stage I and II melanosomes (PubMed:23754390). Colocalizes with CD63 and APOE at exosomes and in intraluminal vesicles within multivesicular endosomes (PubMed:21962903, PubMed:26387950)

#### **Tissue Location**


Normally expressed at low levels in quiescent adult melanocytes but overexpressed by proliferating neonatal melanocytes and during tumor growth. Overexpressed in melanomas. Some expression was found in dysplastic nevi.


### **PMEL17 / GP100 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **PMEL17 / GP100 Antibody - Images**





Western blot analysis of PMEL17 / GP100 expression in human melanoma lysate.