

Phospho-HSF1 (S326) Antibody
Rabbit mAb
Catalog # AP90310**Specification**

Phospho-HSF1 (S326) Antibody - Product Information

Application	WB, IHC, FC, ICC, IP
Primary Accession	Q00613
Clonality	Monoclonal
Other Names	
HSF1; HSTF 1; HSTF1; Heat shock factor protein 1; Heat shock transcription factor 1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	57260 Da

Phospho-HSF1 (S326) Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Phospho-HSF1 (S326)
Description	Heat shock transcription factors (HSF, also designated HSTF) 1 and 2 are involved in this regulation. HSF1 and HSF2 are upregulated by estrogen at both the mRNA and protein level. HSF1 is normally found as a monomer whose transcriptional activity is repressed by constitutive phosphorylation. Upon activation, HSF1 forms trimers, gains DNA binding activity and is translocated to the nucleus. HSF2 activity is associated with differentiation and development, and, like HSF1, binds DNA as a trimer. Both HSF1 and HSF2 are known to be induced by proteasome inhibitors of the ubiquitin pathway.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Phospho-HSF1 (S326) Antibody - Protein Information

Name HSF1 ([HGNC:5224](#))

Synonyms HSTF1

Function

Functions as a stress-inducible and DNA-binding transcription factor that plays a central role in the transcriptional activation of the heat shock response (HSR), leading to the expression of a large class of molecular chaperones, heat shock proteins (HSPs), that protect cells from cellular insult damage (PubMed:[11447121](http://www.uniprot.org/citations/11447121))

(PubMed:[12659875](http://www.uniprot.org/citations/12659875))

(PubMed:[12917326](http://www.uniprot.org/citations/12917326))

(PubMed:[15016915](http://www.uniprot.org/citations/15016915))

(PubMed:[18451878](http://www.uniprot.org/citations/18451878))

(PubMed:[1871105](http://www.uniprot.org/citations/1871105))

(PubMed:[1986252](http://www.uniprot.org/citations/1986252))

(PubMed:[25963659](http://www.uniprot.org/citations/25963659))

(PubMed:[26754925](http://www.uniprot.org/citations/26754925))

(PubMed:[7623826](http://www.uniprot.org/citations/7623826))

(PubMed:[7760831](http://www.uniprot.org/citations/7760831))

(PubMed:[8940068](http://www.uniprot.org/citations/8940068))

(PubMed:[8946918](http://www.uniprot.org/citations/8946918))

(PubMed:[9121459](http://www.uniprot.org/citations/9121459))

(PubMed:[9341107](http://www.uniprot.org/citations/9341107))

(PubMed:[9499401](http://www.uniprot.org/citations/9499401))

(PubMed:[9535852](http://www.uniprot.org/citations/9535852))

(PubMed:[9727490](http://www.uniprot.org/citations/9727490)). In unstressed cells, is present in a HSP90-containing

multichaperone complex that maintains it in a non-DNA-binding inactivated monomeric form

(PubMed:[11583998](http://www.uniprot.org/citations/11583998)),

(PubMed:[16278218](http://www.uniprot.org/citations/16278218)),

(PubMed:[9727490](http://www.uniprot.org/citations/9727490)). Upon exposure to heat and other stress stimuli, undergoes homotrimerization and activates HSP

gene transcription through binding to site-specific heat shock elements (HSEs) present in the

promoter regions of HSP genes (PubMed:[10359787](http://www.uniprot.org/citations/10359787))

(PubMed:[11583998](http://www.uniprot.org/citations/11583998)),

(PubMed:[12659875](http://www.uniprot.org/citations/12659875))

(PubMed:[16278218](http://www.uniprot.org/citations/16278218))

(PubMed:[1871105](http://www.uniprot.org/citations/1871105))

(PubMed:[1986252](http://www.uniprot.org/citations/1986252))

(PubMed:[25963659](http://www.uniprot.org/citations/25963659))

(PubMed:[26754925](http://www.uniprot.org/citations/26754925))

(PubMed:[7623826](http://www.uniprot.org/citations/7623826))

(PubMed:[7935471](http://www.uniprot.org/citations/7935471))

(PubMed:[8455624](http://www.uniprot.org/citations/8455624))

(PubMed:[8940068](http://www.uniprot.org/citations/8940068))

(PubMed:[9499401](http://www.uniprot.org/citations/9499401))

(PubMed:[9727490](http://www.uniprot.org/citations/9727490)). Upon heat shock stress, forms a chromatin-associated complex

with TTC5/STRAP and p300/EP300 to stimulate HSR transcription, therefore increasing cell survival

(PubMed:[18451878](http://www.uniprot.org/citations/18451878)).

Activation is reversible, and during the attenuation and recovery phase period of the HSR, returns

to its unactivated form (PubMed:[11583998](http://www.uniprot.org/citations/11583998))

(PubMed:[16278218](http://www.uniprot.org/citations/16278218)).

Binds to inverted 5'-NGAAN-3' pentamer DNA sequences

(PubMed:[1986252](http://www.uniprot.org/citations/1986252)),

(PubMed:[26727489](http://www.uniprot.org/citations/26727489)).

Binds to chromatin at heat shock gene promoters (PubMed:[11583998](#))

<http://www.uniprot.org/citations/25963659> target="_blank">25963659). Activates transcription of transcription factor FOXR1 which in turn activates transcription of the heat shock chaperones HSPA1A and HSPA6 and the antioxidant NADPH-dependent reductase DHRS2 (PubMed:34723967). Also serves several other functions independently of its transcriptional activity. Involved in the repression of Ras-induced transcriptional activation of the c-fos gene in heat-stressed cells (PubMed:9341107). Positively regulates pre-mRNA 3'-end processing and polyadenylation of HSP70 mRNA upon heat-stressed cells in a symplekin (SYMPK)-dependent manner (PubMed:14707147). Plays a role in nuclear export of stress- induced HSP70 mRNA (PubMed:17897941). Plays a role in the regulation of mitotic progression (PubMed:18794143). Also plays a role as a negative regulator of non-homologous end joining (NHEJ) repair activity in a DNA damage-dependent manner (PubMed:26359349). Involved in stress-induced cancer cell proliferation in a IER5-dependent manner (PubMed:26754925).


Cellular Location

Nucleus. Cytoplasm. Nucleus, nucleoplasm. Cytoplasm, perinuclear region. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Chromosome, centromere, kinetochore Note=The monomeric form is cytoplasmic in unstressed cells (PubMed:26159920, PubMed:8455624). Predominantly nuclear protein in both unstressed and heat shocked cells (PubMed:10359787, PubMed:10413683). Translocates in the nucleus upon heat shock (PubMed:8455624). Nucleocytoplasmic shuttling protein (PubMed:26159920). Colocalizes with IER5 in the nucleus (PubMed:27354066). Colocalizes with BAG3 to the nucleus upon heat stress (PubMed:26159920, PubMed:8455624). Localizes in subnuclear granules called nuclear stress bodies (nSBs) upon heat shock (PubMed:10359787, PubMed:10747973, PubMed:11447121, PubMed:11514557, PubMed:19229036, PubMed:24581496, PubMed:25963659). Colocalizes with SYMPK and SUMO1 in nSBs upon heat shock (PubMed:10359787, PubMed:11447121, PubMed:11514557, PubMed:12665592, PubMed:14707147) Colocalizes with PRKACA/PKA in the nucleus and nSBs upon heat shock (PubMed:21085490). Relocalizes from the nucleus to the cytoplasm during the attenuation and recovery phase period of the heat shock response (PubMed:26159920). Translocates in the cytoplasm in a YWHAE- and XPO1/CRM1-dependent manner (PubMed:12917326). Together with histone H2AX, redistributed in discrete nuclear DNA damage-induced foci after ionizing radiation (IR) (PubMed:26359349). Colocalizes with calcium-responsive transactivator SS18L1 at kinetochore region on the mitotic chromosomes (PubMed:18794143). Colocalizes with gamma tubulin at centrosome (PubMed:18794143). Localizes at spindle pole in metaphase (PubMed:18794143). Colocalizes with PLK1 at spindle poles during prometaphase (PubMed:18794143).

Phospho-HSF1 (S326) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-HSF1 (S326) Antibody - Images

Western blot analysis of Phospho-HSF1 (S326) expression in HeLa cell lysates treated with heat.