

## **TrkA Antibody**

Rabbit mAb Catalog # AP90370

## **Specification**

## **TrkA Antibody - Product Information**

Application WB, IHC, FC, ICC, IP

Primary Accession
Reactivity
Rat
Clonality
Monoclonal

**Other Names** 

NTRK1; MTC; TRK; TRK1; TRKA; Trk-A; p140-TrkA;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 87497 Da

# **TrkA Antibody - Additional Information**

Dilution WB~~1:1000

IHC~~1:100~500 FC~~1:10~50 ICC~~N/A IP~~N/A

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

**TrkA** 

Description The family of Trk receptor tyrosine kinases

consists of TrkA, TrkB, and TrkC. While the sequence of these family members is highly conserved, they are activated by different neurotrophins: TrkA by NGF, TrkB

by BDNF or NT4, and TrkC by NT3. Neurotrophin signaling through these receptors regulates a number of physiological processes, such as cell

survival, proliferation, neural

development, and axon and dendrite

growth and patterning.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

## **TrkA Antibody - Protein Information**

Name NTRK1



#### **Function**

Receptor tyrosine kinase involved in the development and the maturation of the central and peripheral nervous systems through regulation of proliferation, differentiation and survival of sympathetic and nervous neurons. High affinity receptor for NGF which is its primary ligand (PubMed:<a href="http://www.uniprot.org/citations/1281417" target=" blank">1281417</a>, PubMed:<a href="http://www.uniprot.org/citations/15488758" target=" blank">15488758</a>, PubMed: <a href="http://www.uniprot.org/citations/17196528" target="blank">17196528</a>, PubMed:<a href="http://www.uniprot.org/citations/1849459" target=" blank">1849459</a>, PubMed:<a href="http://www.uniprot.org/citations/1850821" target="\_blank">1850821</a>, PubMed:<a href="http://www.uniprot.org/citations/22649032" target="\_blank">22649032</a>, PubMed:<a href="http://www.uniprot.org/citations/22649032" target="\_blank">22649032</a>, PubMed:<a href="http://www.uniprot.org/citations/27445338" target=" blank">27445338</a>, PubMed:<a href="http://www.uniprot.org/citations/8325889" target=" blank">8325889</a>). Can also bind and be activated by NTF3/neurotrophin-3. However, NTF3 only supports axonal extension through NTRK1 but has no effect on neuron survival (By similarity). Upon dimeric NGF ligand-binding, undergoes homodimerization, autophosphorylation and activation (PubMed: <a href="http://www.uniprot.org/citations/1281417" target=" blank">1281417</a>). Recruits, phosphorylates and/or activates several downstream effectors including SHC1, FRS2, SH2B1, SH2B2 and PLCG1 that regulate distinct overlapping signaling cascades driving cell survival and differentiation. Through SHC1 and FRS2 activates a GRB2-Ras-MAPK cascade that regulates cell differentiation and survival. Through PLCG1 controls NF-Kappa-B activation and the transcription of genes involved in cell survival. Through SHC1 and SH2B1 controls a Ras-PI3 kinase-AKT1 signaling cascade that is also regulating survival. In absence of ligand and activation, may promote cell death, making the survival of neurons dependent on trophic factors.

#### **Cellular Location**

Cell membrane; Single-pass type I membrane protein. Early endosome membrane {ECO:0000250|UniProtKB:P35739}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P35739}. Late endosome membrane {ECO:0000250|UniProtKB:P35739}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P35739}. Recycling endosome membrane {ECO:0000250|UniProtKB:P35739}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P35739}. Note=Rapidly internalized after NGF binding (PubMed:1281417). Internalized to endosomes upon binding of NGF or NTF3 and further transported to the cell body via a retrograde axonal transport. Localized at cell membrane and early endosomes before nerve growth factor (NGF) stimulation. Recruited to late endosomes after NGF stimulation. Colocalized with RAPGEF2 at late endosomes {ECO:0000250|UniProtKB:P35739, ECO:0000269|PubMed:1281417}

### **Tissue Location**

Isoform TrkA-I is found in most non-neuronal tissues. Isoform TrkA-II is primarily expressed in neuronal cells TrkA-III is specifically expressed by pluripotent neural stem and neural crest progenitors.

### **TrkA Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture



# **TrkA Antibody - Images**

