

Alkaline Phosphatase Antibody Rabbit mAb Catalog # AP90448

### **Specification**

# **Alkaline Phosphatase Antibody - Product Information**

ApplicationWB, IHC, FC, ICC, IPPrimary AccessionP05186ReactivityRatClonalityMonoclonalOther NamesAlkaline phosphatase, tissue-nonspecific isozyme; AP-TNAP; TNSALP; Alkaline phosphataseliver/bone/kidney isozyme; ALPL;

lsotype	Rabbit IgG
Host	Rabbit
Calculated MW	57305 Da

# **Alkaline Phosphatase Antibody - Additional Information**

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Alkaline Phosphatase
Description	There are at least four distinct but related alkaline phosphatases: intestinal, placental, placental-like, and liver/bone/kidney (tissue non-specific). The first three are located together on chromosome 2, while the tissue non-specific form is located on chromosome 1. The product of this gene is a membrane bound glycosylated enzyme that is not expressed in any particular tissue and is, therefore, referred to as the tissue-nonspecific form of the enzyme.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

# Alkaline Phosphatase Antibody - Protein Information

Name ALPL {ECO:0000303|PubMed:8406453, ECO:0000312|HGNC:HGNC:438}



#### Function

Alkaline phosphatase that metabolizes various phosphate compounds and plays a key role in skeletal mineralization and adaptive thermogenesis (PubMed:<a

href="http://www.uniprot.org/citations/12162492" target="\_blank">12162492</a>, PubMed:<a href="http://www.uniprot.org/citations/23688511" target=" blank">23688511</a>, PubMed:<a href="http://www.uniprot.org/citations/25982064" target=" blank">25982064</a>). Has broad substrate specificity and can hydrolyze a considerable variety of compounds: however, only a few substrates, such as diphosphate (inorganic pyrophosphate; PPi), pyridoxal 5'-phosphate (PLP) and N- phosphocreatine are natural substrates (PubMed: <a href="http://www.uniprot.org/citations/12162492" target="\_blank">12162492</a>, PubMed:<a href="http://www.uniprot.org/citations/2220817" target=" blank">2220817</a>). Plays an essential role in skeletal and dental mineralization via its ability to hydrolyze extracellular diphosphate, a potent mineralization inhibitor, to phosphate: it thereby promotes hydroxyapatite crystal formation and increases inorganic phosphate concentration (PubMed:<a href="http://www.uniprot.org/citations/23688511" target=" blank">23688511</a>, PubMed:<a href="http://www.uniprot.org/citations/25982064" target=" blank">25982064</a>). Acts in a non- redundant manner with PHOSPHO1 in skeletal mineralization: while PHOSPHO1 mediates the initiation of hydroxyapatite crystallization in the matrix vesicles (MVs), ALPL/TNAP catalyzes the spread of hydroxyapatite crystallization in the extracellular matrix (By similarity). Also promotes dephosphorylation of osteopontin (SSP1), an inhibitor of hydroxyapatite crystallization in its phosphorylated state; it is however unclear whether ALPL/TNAP mediates SSP1 dephosphorylation via a direct or indirect manner (By similarity). Catalyzes dephosphorylation of PLP to pyridoxal (PL), the transportable form of vitamin B6, in order to provide a sufficient amount of PLP in the brain, an essential cofactor for enzymes catalyzing the synthesis of diverse neurotransmitters (PubMed:<a href="http://www.uniprot.org/citations/20049532" target="\_blank">20049532</a>, PubMed:<a href="http://www.uniprot.org/citations/2220817" target=" blank">2220817</a>). Additionally, also able to mediate ATP degradation in a stepwise manner to adenosine, thereby regulating the availability of ligands for purinergic receptors (By similarity). Also capable of dephosphorylating microbial products, such as lipopolysaccharides (LPS) as well as other phosphorylated small-molecules, such as poly-inosine:cytosine (poly I:C) (PubMed:<a href="http://www.uniprot.org/citations/28448526" target=" blank">28448526</a>). Acts as a key regulator of adaptive thermogenesis as part of the futile creatine cycle: localizes to the mitochondria of thermogenic fat cells and acts by mediating hydrolysis of N-phosphocreatine to initiate a futile cycle of creatine dephosphorylation and phosphorylation (By similarity). During the futile creatine cycle, creatine and N-phosphocreatine are in a futile cycle, which dissipates the high energy charge of N-phosphocreatine as heat without performing any mechanical or chemical work (By similarity).

#### **Cellular Location**

Cell membrane; Lipid-anchor, GPI-anchor Extracellular vesicle membrane {ECO:000250|UniProtKB:P09242}; Lipid- anchor, GPI-anchor {ECO:000250|UniProtKB:P09242}. Mitochondrion membrane {ECO:000250|UniProtKB:P09242}; Lipid-anchor, GPI-anchor {ECO:0000250|UniProtKB:P09242}. Mitochondrion intermembrane space {ECO:0000250|UniProtKB:P09242}. Note=Localizes to special class of extracellular vesicles, named matrix vesicles (MVs), which are released by osteogenic cells. Localizes to the mitochondria of thermogenic fat cells: tethered to mitochondrial membranes via a GPI-anchor and probably resides in the mitochondrion intermembrane space {ECO:000250|UniProtKB:P09242}

#### Alkaline Phosphatase Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot



- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

# **Alkaline Phosphatase Antibody - Images**



Western blot analysis of Alkaline Phosphatase expression in (1)HepG2 cell lysate; (2)JAR cell lysate.