

# Fos B Antibody

Rabbit mAb Catalog # AP90483

# **Specification**

# Fos B Antibody - Product Information

Application WB, IHC, ICC, IP

Primary Accession P53539
Reactivity Rat

Clonality Monoclonal

**Other Names** 

G0/G1 switch regulatory protein 3, G0S3, Protein fosB; GOSB; Oncogene FOSB; Activator protein 1;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 35928 Da

# Fos B Antibody - Additional Information

Dilution WB~~1:1000

IHC~~1:100~500

ICC~~N/A IP~~N/A

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

Fos B

Description FosB interacts with Jun proteins enhancing

their DNA binding activity. The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex

AP-1.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline ,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

# Fos B Antibody - Protein Information

**Name FOSB** 

Synonyms G0S3

## **Function**

Heterodimerizes with proteins of the JUN family to form an AP-1 transcription factor complex, thereby enhancing their DNA binding activity to gene promoters containing an AP-1 consensus



sequence 5'- TGA[GC]TCA-3' and enhancing their transcriptional activity (PubMed:<a href="http://www.uniprot.org/citations/12618758" target="\_blank">12618758</a>, PubMed:<a href="http://www.uniprot.org/citations/28981703" target="\_blank">28981703</a>). As part of the AP-1 complex, facilitates enhancer selection together with cell-type-specific transcription factors by collaboratively binding to nucleosomal enhancers and recruiting the SWI/SNF (BAF) chromatin remodeling complex to establish accessible chromatin (By similarity). Together with JUN, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/12618758" target="\_blank">12618758</a>(a>). Exhibits transactivation activity in vitro (By similarity). Involved in the display of nurturing behavior towards newborns (By similarity). May play a role in neurogenesis in the hippocampus and in learning and memory-related tasks by regulating the expression of various genes involved in neurogenesis, depression and epilepsy (By similarity). Implicated in behavioral responses related to morphine reward and spatial memory (By similarity).

### **Cellular Location**

Nucleus {ECO:0000250|UniProtKB:P13346}.

#### **Tissue Location**


[Isoform 11]: Expressed in the nucleus accumbens of the striatum (at protein level).

# Fos B Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# Fos B Antibody - Images



Western blot analysis of FosB expression in C6 cell lysate treated with serum.