

**PDGFR alpha Antibody**  
**Rabbit mAb**  
**Catalog # AP91145****Specification**

---

**PDGFR alpha Antibody - Product Information**

|  |                        |
|--|------------------------|
| Application                                | WB                     |
| Primary Accession                          | <a href="#">P16234</a> |
| Reactivity                                 | Rat                    |
| Clonality                                  | Monoclonal             |
| <b>Other Names</b>                         |                        |
| PDGF-R-alpha; PDGFR-alpha; CD140a antigen; |                        |
| Isotype                                    | Rabbit IgG             |
| Host                                       | Rabbit                 |
| Calculated MW                              | 122670 Da              |

**PDGFR alpha Antibody - Additional Information**

|                              |  |
|------------------------------|--|
| Dilution                     | WB~~1:1000   |
| Purification                 | Affinity-chromatography  |
| Immunogen                    | A synthesized peptide derived from human PDGFR alpha   |
| Description                  | Platelet derived growth factor (PDGF) family proteins exist as several disulphide-bonded, dimeric isoforms (PDGF AA, PDGF AB, PDGF BB, PDGF CC, and PDGF DD) that bind in a specific pattern to two closely related receptor tyrosine kinases, PDGF receptor $\alpha$ (PDGFR $\alpha$ ) and PDGF receptor $\beta$ (PDGFR $\beta$ ). Receptor that binds both PDGFA and PDGFB and has a tyrosine-protein kinase activity. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.  |

**PDGFR alpha Antibody - Protein Information****Name** PDGFRA**Synonyms** PDGFR2, RHEPDGFRA**Function**

Tyrosine-protein kinase that acts as a cell-surface receptor for PDGFA, PDGFB and PDGFC and plays an essential role in the regulation of embryonic development, cell proliferation, survival and chemotaxis. Depending on the context, promotes or inhibits cell proliferation and cell migration.

Plays an important role in the differentiation of bone marrow-derived mesenchymal stem cells. Required for normal skeleton development and cephalic closure during embryonic development. Required for normal development of the mucosa lining the gastrointestinal tract, and for recruitment of mesenchymal cells and normal development of intestinal villi. Plays a role in cell migration and chemotaxis in wound healing. Plays a role in platelet activation, secretion of agonists from platelet granules, and in thrombin-induced platelet aggregation. Binding of its cognate ligands - homodimeric PDGFA, homodimeric PDGFB, heterodimers formed by PDGFA and PDGFB or homodimeric PDGFC - leads to the activation of several signaling cascades; the response depends on the nature of the bound ligand and is modulated by the formation of heterodimers between PDGFRA and PDGFRB. Phosphorylates PIK3R1, PLCG1, and PTPN11. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate, mobilization of cytosolic  $\text{Ca}^{2+}$  and the activation of protein kinase C. Phosphorylates PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase, and thereby mediates activation of the AKT1 signaling pathway. Mediates activation of HRAS and of the MAP kinases MAPK1/ERK2 and/or MAPK3/ERK1. Promotes activation of STAT family members STAT1, STAT3 and STAT5A and/or STAT5B. Receptor signaling is down-regulated by protein phosphatases that dephosphorylate the receptor and its down-stream effectors, and by rapid internalization of the activated receptor.

#### **Cellular Location**

Cell membrane; Single-pass type I membrane protein. Cell projection, cilium {ECO:0000250|UniProtKB:P26618}. Golgi apparatus {ECO:0000250|UniProtKB:P26618}

#### **Tissue Location**

Detected in platelets (at protein level). Widely expressed. Detected in brain, fibroblasts, smooth muscle, heart, and embryo. Expressed in primary and metastatic colon tumors and in normal colon tissue.


### **PDGFR alpha Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **PDGFR alpha Antibody - Images**





Western blot analysis of PDGFR expression in SH-SY5Y cell lysate.