

**NeuroD1 Antibody**  
**Rabbit mAb**  
**Catalog # AP91211****Specification**

---

**NeuroD1 Antibody - Product Information**

Application	WB, IP
Primary Accession	<a href="#">Q13562</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
BETA2; BHF1; bHLHa3; MODY6; NDF1; NEUROD; NeuroD1; Neurogenic;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	39920 Da

**NeuroD1 Antibody - Additional Information**

Dilution	WB~~1:1000 IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human NeuroD1
Description	Differentiation factor required for dendrite morphogenesis and maintenance in the cerebellar cortex. Transcriptional activator. Binds to the insulin gene E-box.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**NeuroD1 Antibody - Protein Information****Name** NEUROD1**Synonyms** BHLHA3, NEUROD**Function**

Acts as a transcriptional activator: mediates transcriptional activation by binding to E box-containing promoter consensus core sequences 5'-CANNTG-3'. Associates with the p300/CBP transcription coactivator complex to stimulate transcription of the secretin gene as well as the gene encoding the cyclin-dependent kinase inhibitor CDKN1A. Contributes to the regulation of several cell differentiation pathways, like those that promote the formation of early retinal ganglion cells, inner ear sensory neurons, granule cells forming either the cerebellum or the dentate gyrus cell layer of the hippocampus, endocrine islet cells of the pancreas and enteroendocrine cells of the small intestine. Together with PAX6 or SIX3, is required for the

regulation of amacrine cell fate specification. Also required for dendrite morphogenesis and maintenance in the cerebellar cortex. Associates with chromatin to enhancer regulatory elements in genes encoding key transcriptional regulators of neurogenesis (By similarity).

#### Cellular Location


Cytoplasm. Nucleus {ECO:0000255|PROSITE-ProRule:PRU00981, ECO:0000269|PubMed:14752053} Note=In pancreatic islet cells, shuttles to the nucleus in response to glucose stimulation (By similarity). Colocalizes with NR0B2 in the nucleus.

#### NeuroD1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### NeuroD1 Antibody - Images



Western blot analysis of NeuroD1 expression in Y79 cell lysate.