

ROCK2 Antibody

Rabbit mAb Catalog # AP91415

# Specification

# **ROCK2 Antibody - Product Information**

ApplicationWB, ICCPrimary Accession075116ReactivityRatClonalityMonoclonalOther Names9164 ROCK 2; Rho kinase 2; ROCK 2; Rock II; Rock2m; ROK alpha; ROKalpha;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	<b>160900 Da</b>

## **ROCK2** Antibody - Additional Information

Dilution

Purification Immunogen

Description

Storage Condition and Buffer

WB~~1:1000 ICC~~N/A Affinity-chromatography A synthesized peptide derived from human ROCK2 Regulates the assembly of the actin cytoskeleton. Promotes formation of stress fibers and of focal adhesion complexes. Plays a role in smooth muscle contraction. Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## **ROCK2 Antibody - Protein Information**

Name ROCK2

Synonyms KIAA0619

Function

Protein kinase which is a key regulator of actin cytoskeleton and cell polarity. Involved in regulation of smooth muscle contraction, actin cytoskeleton organization, stress fiber and focal adhesion formation, neurite retraction, cell adhesion and motility via phosphorylation of ADD1, BRCA2, CNN1, EZR, DPYSL2, EP300, MSN, MYL9/MLC2, NPM1, RDX, PPP1R12A and VIM. Phosphorylates SORL1 and IRF4. Acts as a negative regulator of VEGF-induced angiogenic endothelial cell activation. Positively regulates the activation of p42/MAPK1- p44/MAPK3 and of p90RSK/RPS6KA1 during myogenic differentiation. Plays an important role in the timely initiation of centrosome duplication. Inhibits keratinocyte terminal differentiation. May regulate closure of the



eyelids and ventral body wall through organization of actomyosin bundles. Plays a critical role in the regulation of spine and synaptic properties in the hippocampus. Plays an important role in generating the circadian rhythm of the aortic myofilament Ca(2+) sensitivity and vascular contractility by modulating the myosin light chain phosphorylation.

#### **Cellular Location**

Cytoplasm. Cell membrane; Peripheral membrane protein. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Cytoplasmic, and associated with actin microfilaments and the plasma membrane.

#### **Tissue Location**


Expressed in the brain (at protein level).

### **ROCK2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

### **ROCK2 Antibody - Images**



Western blot analysis of ROCK2 expression in (1) Hela cell lysate; (2) RAW264.7 cell lysate.