

WIPI1 Antibody

Rabbit mAb Catalog # AP91665

Specification

WIPI1 Antibody - Product Information

Application WB, IHC
Primary Accession Q5MNZ9
Reactivity Rat
Clonality Monoclonal

Other Names

ATG18; ATG18A; WIPI 1; WIPI-1; WIPI49;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 48673 Da

WIPI1 Antibody - Additional Information

Dilution WB~~1:1000

IHC~~1:100~500

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

WIPI1

Description May play a role in autophagy. May regulate

the trafficking of proteins involved in the mannose-6-phosphate receptor (MPR)

recycling pathway.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline ,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

WIPI1 Antibody - Protein Information

Name WIPI1

Synonyms WIPI49

Function

Component of the autophagy machinery that controls the major intracellular degradation process by which cytoplasmic materials are packaged into autophagosomes and delivered to lysosomes for degradation (PubMed:<a href="http://www.uniprot.org/citations/15602573"

 $target="_blank">15602573, PubMed:20114074, PubMed:20114074, PubMed:20114074, PubMed:<a$

target="_blank">20114074, PubMed:20484055, PubMed:<a href="http://www.uniprot.org/citations/20639694"

target="_blank">20639694, PubMed:<a href="http://www.uniprot.org/citations/23088497"

target="blank">23088497, PubMed:<a href="http://www.uniprot.org/citations/28561066"



target=" blank">28561066, PubMed:31271352). Plays an important role in starvation- and calcium-mediated autophagy, as well as in mitophagy (PubMed: 28561066). Functions downstream of the ULK1 and PI3- kinases that produce phosphatidylinositol 3-phosphate (PtdIns3P) on membranes of the endoplasmic reticulum once activated (PubMed:28561066). Binds phosphatidylinositol 3-phosphate (PtdIns3P), and maybe other phosphoinositides including PtdIns3,5P2 and PtdIns5P, and is recruited to phagophore assembly sites at the endoplasmic reticulum membranes (PubMed: <a $href="http://www.uniprot.org/citations/28561066" target="_blank">28561066, PubMed:31271352, PubMed:31271352, PubMed:33499712). There, it assists WIPI2 in the recruitment of ATG12- ATG5-ATG16L1, a complex that directly controls the elongation of the nascent autophagosomal membrane (PubMed: 28561066). Together with WDR45/WIPI4, promotes ATG2 (ATG2A or ATG2B)-mediated lipid transfer by enhancing ATG2-association with phosphatidylinositol 3-monophosphate (PI3P)-containing membranes

Involved in xenophagy of Staphylococcus aureus (PubMed:22829830). Invading S.aureus cells become entrapped in autophagosome-like WIPI1 positive vesicles targeted for lysosomal degradation (PubMed: 22829830). Also plays a distinct role in controlling the transcription of melanogenic enzymes and melanosome maturation, a process that is distinct from starvation-induced autophagy (PubMed:21317285). May also regulate the trafficking of proteins involved in the mannose-6-phosphate receptor (MPR) recycling pathway (PubMed:<a

(PubMed:31271352).

href="http://www.uniprot.org/citations/15020712" target=" blank">15020712).

Cellular Location

Golgi apparatus, trans-Golgi network. Endosome. Cytoplasmic vesicle, clathrin-coated vesicle. Preautophagosomal structure membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Note=Trans elements of the Golgi and peripheral endosomes. Dynamically cycles through these compartments and is susceptible to conditions that modulate membrane flux. Enriched in clathrin-coated vesicles. Upon starvation-induced autophagy, accumulates at subcellular structures in the cytoplasm: enlarged vesicular and lasso-like structures, and large cup-shaped structures predominantly around the nucleus. Recruitment to autophagic membranes is controlled by MTMR14. Labile microtubules specifically recruit markers of autophagosome formation like WIPI1, whereas mature autophagosomes may bind to stable microtubules

Tissue Location

Ubiquitously expressed. Highly expressed in skeletal muscle, heart, testis, pancreas and placenta. Highly expressed in G361, Sk-mel-28, Sk-mel-13, WM852 and WM451 cells. Up-regulated in a variety of tumor tissues.

WIPI1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation



Tel: 858.875.1900 Fax: 858.875.1999

- Flow Cytomety
- Cell Culture

WIPI1 Antibody - Images



Western blot analysis of WIPI1 expression in A375 cell lysate.