

FCGR1B Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP9197B

Specification

FCGR1B Antibody (C-term) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Antigen Region WB, FC, IHC-P-Leica,E <u>O92637</u> <u>A6NKC4</u>, <u>P12314</u> Human Rabbit Polyclonal Rabbit IgG 248-276

FCGR1B Antibody (C-term) - Additional Information

Other Names

High affinity immunoglobulin gamma Fc receptor IB, IgG Fc receptor IB, Fc-gamma RIB, FcRIB, hFcgammaRIB, FCGR1B, IGFRB

Target/Specificity

This FCGR1B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 248-276 amino acids from the C-terminal region of human FCGR1B.

Dilution WB~~1:500 FC~~1:10~50 IHC-P-Leica~~1:100 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FCGR1B Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

FCGR1B Antibody (C-term) - Protein Information

Name FCGR1BP (<u>HGNC:3614</u>)

Synonyms FCGR1B, IGFRB



Function May bind to the Fc region of immunoglobulins gamma with a low affinity compared to FCGR1A. May function in the humoral immune response.

Cellular Location


Cell membrane; Single-pass type I membrane protein

FCGR1B Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

FCGR1B Antibody (C-term) - Images




All lanes : Anti-FCGR1B Antibody (C-term) at 1:500 dilution Lane 1: HL-60 whole cell lysate Lane 2: Hela whole cell lysate Lane 3: K562 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 33 kDa Blocking/Dilution buffer: 5% NFDM/TBST.





All lanes : Anti-FCGR1B Antibody (C-term) at 1:500 dilution Lane 1: CCRF-CEM whole cell lysate Lane 2: THP-1 whole cell lysate Lane 3: Hela whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 33 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Immunohistochemical analysis of AP9197b on paraffin-embedded human tonsil tissue was performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 15min at room temperature. Leica Bond Polymer Refine Detection was used as the secondary antibody.





Immunohistochemical analysis of AP9197b on paraffin-embedded human appendix tissue was performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 15min at room temperature. Leica Bond Polymer Refine Detection was used as the secondary antibody.



FCGR1B Antibody (C-term) (Cat. #AP9197b) flow cytometric analysis of CEM cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

FCGR1B Antibody (C-term) - Background

FCGR1B may bind to the Fc region of immunoglobulins gamma with a low affinity compared to FCGR1A. This protein may function in the humoral immune response.

FCGR1B Antibody (C-term) - References

Thomas,G., et.al., Nat. Genet. 41 (5), 579-584 (2009) Kuwano,Y., et.al., Arch. Dermatol. Res. 298 (10), 493-498 (2007)