

AKAP 95 Antibody
Rabbit mAb
Catalog # AP92911**Specification**

AKAP 95 Antibody - Product Information

Application	WB, IHC, ICC
Primary Accession	O43823
Reactivity	Rat
Clonality	Monoclonal
Other Names	
Akap8; AKAP95;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	76108 Da

AKAP 95 Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human AKAP 95
Description	Anchoring protein that mediates the subcellular compartmentation of cAMP-dependent protein kinase (PKA type II).
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

AKAP 95 Antibody - Protein Information**Name** AKAP8**Synonyms** AKAP95**Function**

Anchoring protein that mediates the subcellular compartmentation of cAMP-dependent protein kinase (PKA type II) (PubMed:9473338). Acts as an anchor for a PKA-signaling complex onto mitotic chromosomes, which is required for maintenance of chromosomes in a condensed form throughout mitosis. Recruits condensin complex subunit NCAPD2 to chromosomes required for chromatin condensation; the function appears to be independent from PKA-anchoring (PubMed:10601332, PubMed:10601332).

[>10791967, PubMed:11964380\). May help to deliver cyclin D/E to CDK4 to facilitate cell cycle progression \(PubMed:14641107\). Required for cell cycle G2/M transition and histone deacetylation during mitosis. In mitotic cells recruits HDAC3 to the vicinity of chromatin leading to deacetylation and subsequent phosphorylation at 'Ser-10' of histone H3; in this function may act redundantly with AKAP8L \(PubMed:16980585\). Involved in nuclear retention of RPS6KA1 upon ERK activation thus inducing cell proliferation \(PubMed:22130794\). May be involved in regulation of DNA replication by acting as scaffold for MCM2 \(PubMed:12740381\). Enhances HMT activity of the KMT2 family MLL4/WBP7 complex and is involved in transcriptional regulation. In a teratocarcinoma cell line is involved in retinoic acid-mediated induction of developmental genes implicating H3 'Lys-4' methylation \(PubMed:23995757\). May be involved in recruitment of active CASP3 to the nucleus in apoptotic cells \(PubMed:16227597\). May act as a carrier protein of GJA1 for its transport to the nucleus \(PubMed:26880274\). May play a repressive role in the regulation of rDNA transcription. Preferentially binds GC-rich DNA in vitro. In cells, associates with ribosomal RNA \(rRNA\) chromatin, preferentially with rRNA promoter and transcribed regions \(PubMed:26683827\). Involved in modulation of Toll- like receptor signaling. Required for the cAMP-dependent suppression of TNF-alpha in early stages of LPS-induced macrophage activation; the function probably implicates targeting of PKA to NFKB1 \(By similarity\).](http://www.uniprot.org/citations/10791967)

Cellular Location

Nucleus. Nucleus matrix. Nucleus, nucleolus. Cytoplasm {ECO:0000250|UniProtKB:Q9DBR0}. Note=Associated with the nuclear matrix in interphase and redistributes mostly to chromatin at mitosis However, mitotic chromatin localization has been questioned. Upon nuclear reassembly at the end of mitosis, is sequestered into the daughter nuclei where it re-acquires an interphase distribution Exhibits partial localization to the nucleolus in interphase, where it colocalizes with UBTF/UBF, suggesting localization to the fibrillary center and/or to the dense fibrillary component. Colocalizes with GJA1 at the nuclear membrane specifically during cell cycle G1/S phase

Tissue Location


Highly expressed in heart, liver, skeletal muscle, kidney and pancreas. Expressed in mature dendritic cells


AKAP 95 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

AKAP 95 Antibody - Images





Western blot analysis of AKAP 95 expression in HepG2 cell lysate.