

Specification

Component of the Wnt-Fzd-LRP5-LRP6 complex that triggers beta-catenin signaling through inducing aggregation of receptor-ligand complexes into ribosome-sized signalosomes (PubMed:[11357136](http://www.uniprot.org/citations/11357136), PubMed:[11448771](http://www.uniprot.org/citations/11448771), PubMed:[15778503](http://www.uniprot.org/citations/15778503), PubMed:[16341017](http://www.uniprot.org/citations/16341017), PubMed:[16513652](http://www.uniprot.org/citations/16513652), PubMed:[17326769](http://www.uniprot.org/citations/17326769), PubMed:[17400545](http://www.uniprot.org/citations/17400545))

href="http://www.uniprot.org/citations/19107203" target="_blank">19107203, PubMed:19293931, PubMed:19801552, PubMed:28341812). Cell-surface coreceptor of Wnt/beta- catenin signaling, which plays a pivotal role in bone formation (PubMed:11357136, PubMed:11448771, PubMed:15778503, PubMed:16341017, PubMed:16513652, PubMed:17326769, PubMed:17400545, PubMed:19107203, PubMed:19293931, PubMed:19801552, PubMed:28341812). The Wnt-induced Fzd/LRP6 coreceptor complex recruits DVL1 polymers to the plasma membrane which, in turn, recruits the AXIN1/GSK3B-complex to the cell surface promoting the formation of signalosomes and inhibiting AXIN1/GSK3-mediated phosphorylation and destruction of beta-catenin (PubMed:16513652). Required for posterior patterning of the epiblast during gastrulation (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein Endoplasmic reticulum. Membrane raft. Note=On Wnt signaling, undergoes a cycle of caveolin- or clathrin-mediated endocytosis and plasma membrane location. Released from the endoplasmic reticulum on palmitoylation Mono-ubiquitination retains it in the endoplasmic reticulum in the absence of palmitoylation. On Wnt signaling, phosphorylated, aggregates and colocalizes with AXIN1 and GSK3B at the plasma membrane in LRP6- signalosomes (By similarity). Chaperoned to the plasma membrane by HSP90B1 and MESD (PubMed:23572575). {ECO:0000250|UniProtKB:O88572, ECO:0000269|PubMed:23572575}

Tissue Location


Widely coexpressed with LRP5 during embryogenesis and in adult tissues


Phospho-LRP6 (S1490) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-LRP6 (S1490) Antibody - Images





Western blot analysis of Phospho-LRP6 (S1490) expression in heLa treated with Calyculin cell lysate.