

# Phospho-MEK1 (T292) (13X15) Rabbit Monoclonal Antibody Phospho-MEK1 (T292) (13X15) Rabbit Monoclonal Antibody Catalog # AP93683

# **Specification**

# Phospho-MEK1 (T292) (13X15) Rabbit Monoclonal Antibody - Product Information

Application WB
Primary Accession Q02750
Reactivity Human
Clonality Monoclonal
Calculated MW 43439

# Phospho-MEK1 (T292) (13X15) Rabbit Monoclonal Antibody - Additional Information

**Gene ID 5604** 

#### **Other Names**

Dual specificity mitogen-activated protein kinase kinase 1, MAP kinase kinase 1, MAPKK 1, MKK1, 2.7.12.2, ERK activator kinase 1, MAPK/ERK kinase 1, MEK 1, MAP2K1 (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=6840" target="\_blank">HGNC:6840</a>), MEK1, PRKMK1

Dilution WB~~1:1000

**Storage Conditions** -20°C

# Phospho-MEK1 (T292) (13X15) Rabbit Monoclonal Antibody - Protein Information

Name MAP2K1 (HGNC:6840)

Synonyms MEK1, PRKMK1

#### **Function**

Dual specificity protein kinase which acts as an essential component of the MAP kinase signal transduction pathway. Binding of extracellular ligands such as growth factors, cytokines and hormones to their cell-surface receptors activates RAS and this initiates RAF1 activation. RAF1 then further activates the dual-specificity protein kinases MAP2K1/MEK1 and MAP2K2/MEK2. Both MAP2K1/MEK1 and MAP2K2/MEK2 function specifically in the MAPK/ERK cascade, and catalyze the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in the extracellular signal-regulated kinases MAPK3/ERK1 and MAPK1/ERK2, leading to their activation and further transduction of the signal within the MAPK/ERK cascade. Activates BRAF in a KSR1 or KSR2-dependent manner; by binding to KSR1 or KSR2 releases the inhibitory intramolecular interaction between KSR1 or KSR2 protein kinase and N-terminal domains which promotes KSR1 or KSR2-BRAF dimerization and BRAF activation (PubMed:<a href="http://www.uniprot.org/citations/29433126" target="\_blank">29433126</a>). Depending on the cellular context, this pathway mediates diverse biological functions such as cell growth,



Tel: 858.875.1900 Fax: 858.875.1999

adhesion, survival and differentiation, predominantly through the regulation of transcription, metabolism and cytoskeletal rearrangements. One target of the MAPK/ERK cascade is peroxisome proliferator-activated receptor gamma (PPARG), a nuclear receptor that promotes differentiation and apoptosis. MAP2K1/MEK1 has been shown to export PPARG from the nucleus. The MAPK/ERK cascade is also involved in the regulation of endosomal dynamics, including lysosome processing and endosome cycling through the perinuclear recycling compartment (PNRC), as well as in the fragmentation of the Golgi apparatus during mitosis.

### **Cellular Location**

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, microtubule organizing center, spindle pole body. Cytoplasm. Nucleus Membrane; Peripheral membrane protein. Note=Localizes at centrosomes during prometaphase, midzone during anaphase and midbody during telophase/cytokinesis (PubMed:14737111). Membrane localization is probably regulated by its interaction with KSR1 (PubMed:10409742)

### **Tissue Location**


Widely expressed, with extremely low levels in brain.

# Phospho-MEK1 (T292) (13X15) Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# Phospho-MEK1 (T292) (13X15) Rabbit Monoclonal Antibody - Images



Western blot analysis of MEK5 expression in (1) HeLa cell lysate; (2) HeLa cell treated with Nocodazole.