

# EMP-1 Rabbit pAb

EMP-1 Rabbit pAb Catalog # AP93944

# **Specification**

### **EMP-1** Rabbit pAb - Product Information

Application Reactivity Host Clonality Calculated MW Physical State

Immunogen

Isotype **Purity** 

affinity purified by Protein A

WB, IHC-P, IHC-F, IF

Mouse Rabbit Polyclonal 17 KDa Liquid

KLH conjugated synthetic peptide derived

from mouse EMP-1

laG

Buffer

SUBCELLULAR LOCATION SIMILARITY Important Note 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

Membrane; Multi-pass membrane protein. Belongs to the PMP-22/EMP/MP20 family. This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

#### **Background Descriptions**

Epithelial membrane protein-1 (EMP-1) is a four pass transmembrane protein consisting of 160 amino acids. It is a member of a small family of epithelial membrane proteins. EMP-1 is very similar in structure to its close relative, Peripheral Myelin Protein 22 (PMP22). It is most predominantly expressed in tissues of the gastrointestinal tract but has also been found to be a junctional protein in the liver expressed along the intercellular border. EMP-1 directly interacts with the C-terminus of the P2X7 receptor and may be involved in membrane blebbing. EMP-1 may also be an important regulator in cell communication, signaling, and adhesion. When EMP-1 is overexpressed, cell proliferation is inhibited, S phase is arrested and G1 phase is prolonged in esophogeal cancer cells. EMP-1 may play a role in tumorigenesis and has been identified as a biomarker for gefitinib treatment resistance for patients with lung cancer.

# EMP-1 Rabbit pAb - Additional Information

# Target/Specificity

Most prominently found in the gastrointestinal tract, skin, lung, and brain but not in liver.

#### **Dilution**

- <span class ="dilution WB">WB~~1:1000</span><br \><span class</pre>
- ="dilution\_IHC-P">IHC-P~~N/A</span><br \><span class
- ="dilution IHC-F">IHC-F $\sim$ N/A</span><br/>span class ="dilution IF">IF $\sim$ 1:50 $\sim$ 200</span>

# **Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH



7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


# **EMP-1** Rabbit pAb - Protein Information

# **EMP-1** Rabbit pAb - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# EMP-1 Rabbit pAb - Images




Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EMP-1) Polyclonal Antibody, Unconjugated (AP93944) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (rat small intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide




for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EMP-1) Polyclonal Antibody, Unconjugated (AP93944) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EMP-1) Polyclonal Antibody, Unconjugated (AP93944) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EMP-1) Polyclonal Antibody, Unconjugated (AP93944) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (mouse small intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EMP-1) Polyclonal Antibody, Unconjugated (AP93944) at 1:200 overnight at 4°C,



Tel: 858.875.1900 Fax: 858.875.1999

followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



25 ug total protein per lane of various lysates (see on figure) probed with EMP-1 polyclonal antibody, unconjugated (AP93944) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.

# EMP-1 Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.