

**EMP-1 Rabbit pAb**  
**EMP-1 Rabbit pAb**  
**Catalog # AP93944****Specification****EMP-1 Rabbit pAb - Product Information**

Application	<b>WB, IHC-P, IHC-F, IF</b>
Reactivity	<b>Mouse</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>17 KDa</b>
Physical State	<b>Liquid</b>
Immunogen	<b>KLH conjugated synthetic peptide derived from mouse EMP-1</b>
Isotype	<b>IgG</b>
<b>Purity</b> affinity purified by Protein A	
Buffer	<b>0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.</b>
SUBCELLULAR LOCATION	<b>Membrane; Multi-pass membrane protein.</b>
SIMILARITY	<b>Belongs to the PMP-22/EMP/MP20 family.</b>
Important Note	<b>This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.</b>

**Background Descriptions**

Epithelial membrane protein-1 (EMP-1) is a four pass transmembrane protein consisting of 160 amino acids. It is a member of a small family of epithelial membrane proteins. EMP-1 is very similar in structure to its close relative, Peripheral Myelin Protein 22 (PMP22). It is most predominantly expressed in tissues of the gastrointestinal tract but has also been found to be a junctional protein in the liver expressed along the intercellular border. EMP-1 directly interacts with the C-terminus of the P2X7 receptor and may be involved in membrane blebbing. EMP-1 may also be an important regulator in cell communication, signaling, and adhesion. When EMP-1 is overexpressed, cell proliferation is inhibited, S phase is arrested and G1 phase is prolonged in esophageal cancer cells. EMP-1 may play a role in tumorigenesis and has been identified as a biomarker for gefitinib treatment resistance for patients with lung cancer.

**EMP-1 Rabbit pAb - Additional Information****Target/Specificity**

Most prominently found in the gastrointestinal tract, skin, lung, and brain but not in liver.

**Dilution**

**<span class = "dilution\_WB">WB~~1:1000</span><br \><span class = "dilution\_IHC-P">IHC-P~~N/A</span><br \><span class = "dilution\_IHC-F">IHC-F~~N/A</span><br \><span class = "dilution\_IF">IF~~1:50~200</span>**

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce


**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


**EMP-1 Rabbit pAb - Protein Information****EMP-1 Rabbit pAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**EMP-1 Rabbit pAb - Images**


Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EMP-1) Polyclonal Antibody, Unconjugated (AP93944) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (rat small intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EMP-1) Polyclonal Antibody, Unconjugated (AP93944) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EMP-1) Polyclonal Antibody, Unconjugated (AP93944) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EMP-1) Polyclonal Antibody, Unconjugated (AP93944) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse small intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen

peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EMP-1) Polyclonal Antibody, Unconjugated (AP93944) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



25 ug total protein per lane of various lysates (see on figure) probed with EMP-1 polyclonal antibody, unconjugated (AP93944) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.

#### **EMP-1 Rabbit pAb - Background**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.