

phospho-Calmodulin 1/2/3 (Ser102) Rabbit pAb
phospho-Calmodulin 1/2/3 (Ser102) Rabbit pAb
Catalog # AP94215

Specification

phospho-Calmodulin 1/2/3 (Ser102) Rabbit pAb - Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	PODP23
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	16838

phospho-Calmodulin 1/2/3 (Ser102) Rabbit pAb - Additional Information

Gene ID 801;805;808

Other Names

Calmodulin-1 {ECO:0000312|HGNC:HGNC:1442}, CALM1 {ECO:0000303|PubMed:7925473, ECO:0000312|HGNC:HGNC:1442}

Dilution

IHC-P~~N/A
IHC-F~~N/A
IF~~1:50~200

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

phospho-Calmodulin 1/2/3 (Ser102) Rabbit pAb - Protein Information

Name CALM1 {ECO:0000303|PubMed:7925473, ECO:0000312|HGNC:HGNC:1442}

Function

Calmodulin acts as part of a calcium signal transduction pathway by mediating the control of a large number of enzymes, ion channels, aquaporins and other proteins through calcium-binding (PubMed:[16760425](http://www.uniprot.org/citations/16760425), PubMed:[23893133](http://www.uniprot.org/citations/23893133), PubMed:[26969752](http://www.uniprot.org/citations/26969752), PubMed:[27165696](http://www.uniprot.org/citations/27165696), PubMed:[28890335](http://www.uniprot.org/citations/28890335), PubMed:[31454269](http://www.uniprot.org/citations/31454269), PubMed:[35568036](http://www.uniprot.org/citations/35568036)). Calcium-binding is required for the activation of calmodulin (PubMed:[16760425](http://www.uniprot.org/citations/16760425), PubMed:[16760425](http://www.uniprot.org/citations/16760425), PubMed:[16760425](http://www.uniprot.org/citations/16760425)).

<http://www.uniprot.org/citations/23893133> target="_blank">23893133, PubMed:26969752, PubMed:27165696, PubMed:28890335, PubMed:31454269, PubMed:35568036). Among the enzymes to be stimulated by the calmodulin-calcium complex are a number of protein kinases, such as myosin light-chain kinases and calmodulin-dependent protein kinase type II (CaMK2), and phosphatases (PubMed:16760425, PubMed:23893133, PubMed:26969752, PubMed:27165696, PubMed:28890335, PubMed:31454269, PubMed:35568036). Together with CCP110 and centrin, is involved in a genetic pathway that regulates the centrosome cycle and progression through cytokinesis (PubMed:16760425). Is a regulator of voltage- dependent L-type calcium channels (PubMed:31454269). Mediates calcium- dependent inactivation of CACNA1C (PubMed:26969752). Positively regulates calcium-activated potassium channel activity of KCNN2 (PubMed:27165696). Forms a potassium channel complex with KCNQ1 and regulates electrophysiological activity of the channel via calcium- binding (PubMed:25441029). Acts as a sensor to modulate the endoplasmic reticulum contacts with other organelles mediated by VMP1:ATP2A2 (PubMed:28890335).

Cellular Location

Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cell projection, cilium, flagellum
{ECO:0000250|UniProtKB:P0DP26} Note=Distributed throughout the cell during interphase, but during mitosis becomes dramatically localized to the spindle poles and the spindle microtubules


phospho-Calmodulin 1/2/3 (Ser102) Rabbit pAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

phospho-Calmodulin 1/2/3 (Ser102) Rabbit pAb - Images





Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-CaM I (Ser 102)) Polyclonal Antibody, Unconjugated (AP94215) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

phospho-Calmodulin 1/2/3 (Ser102) Rabbit pAb - Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.