

NDUFS2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9769c

Specification

NDUFS2 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	O75306
Other Accession	O641Y2 , O91WD5 , P17694
Reactivity	Human, Mouse
Predicted	Bovine, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	52546
Antigen Region	286-315

NDUFS2 Antibody (Center) - Additional Information

Gene ID 4720

Other Names

NADH dehydrogenase [ubiquinone] iron-sulfur protein 2, mitochondrial, Complex I-49kD, CI-49kD, NADH-ubiquinone oxidoreductase 49 kDa subunit, NDUFS2

Target/Specificity

This NDUFS2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 286-315 amino acids from the Central region of human NDUFS2.

Dilution

WB~~1:1000

IHC-P~~1:50~100

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NDUFS2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

NDUFS2 Antibody (Center) - Protein Information

Name NDUF52

Function Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which catalyzes electron transfer from NADH through the respiratory chain, using ubiquinone as an electron acceptor (PubMed:[22036843](#), PubMed:[28031252](#), PubMed:[30922174](#)). Essential for the catalytic activity of complex I (PubMed:[22036843](#), PubMed:[30922174](#)). Essential for the assembly of complex I (By similarity). Redox-sensitive, critical component of the oxygen-sensing pathway in the pulmonary vasculature which plays a key role in acute pulmonary oxygen-sensing and hypoxic pulmonary vasoconstriction (PubMed:[30922174](#)). Plays an important role in carotid body sensing of hypoxia (By similarity). Essential for glia-like neural stem and progenitor cell proliferation, differentiation and subsequent oligodendrocyte or neuronal maturation (By similarity).

Cellular Location


Mitochondrion inner membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:Q641Y2}; Matrix side {ECO:0000250|UniProtKB:Q641Y2}

NDUF52 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NDUF52 Antibody (Center) - Images



Western blot analysis of lysates from A431, Hela cell line, mouse brain and rat brain tissue lysate(from left to right), using NDUF52 Antibody (Center)(Cat. #AP9769c). AP9769c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



NDUFS2 Antibody (Center) (Cat. #AP9769c) IHC analysis in formalin fixed and paraffin embedded hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the NDUFS2 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

NDUFS2 Antibody (Center) - Background

NDUFS2 is a core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (complex I). Mammalian mitochondrial complex I is composed of at least 43 different subunits, 7 of which are encoded by the mitochondrial genome, and the rest are the products of nuclear genes. The iron-sulfur protein fraction of complex I is made up of 7 subunits, including this gene product. Complex I catalyzes the NADH oxidation with concomitant ubiquinone reduction and proton ejection out of the mitochondria.

NDUFS2 Antibody (Center) - References

Saada, A., et al. Am. J. Hum. Genet. 84(6):718-727(2009) Wang, L., et al. Cancer Epidemiol. Biomarkers Prev. 17(12):3558-3566(2008) Starr, J.M., et al. Mech. Ageing Dev. 129(12):745-751(2008) Ban, M., et al. PLoS ONE 3 (8), E2891 (2008) Triepels, R.H., et al. J. Biol. Chem. 276(12):8892-8897(2001)