

FCRL4 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP9847b

Specification

FCRL4 Antibody (C-term) - Product Information

Application FC, WB, E
Primary Accession Q96PJ5
Reactivity Human
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Antigen Region 480-507

FCRL4 Antibody (C-term) - Additional Information

Gene ID 83417

Other Names

Fc receptor-like protein 4, FcR-like protein 4, FcRL4, Fc receptor homolog 4, FcRH4, IFGP family protein 2, hIFGP2, Immune receptor translocation-associated protein 1, CD307d, FCRL4, FCRH4, IFGP2, IRTA1

Target/Specificity

This FCRL4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 480-507 amino acids from the C-terminal region of human FCRL4.

Dilution

FC~~1:10~50 WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FCRL4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

FCRL4 Antibody (C-term) - Protein Information

Name FCRL4





Synonyms FCRH4, IFGP2, IRTA1

Function May function as an inhibitor of the B-cell receptor signaling. May function in the B-cell-mediated immune response.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location


Specifically expressed by memory and monocytoid B- cells which populate spleen and lymph nodes. Preferentially expressed in memory B-cells associated with mucosal tissue (at protein level)

FCRL4 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

FCRL4 Antibody (C-term) - Images




All lanes: Anti-FCRL4 Antibody (C-term) at 1:2000 dilution Lane 1: 293T/17 whole cell lysate Lane 2: Ramos whole cell lysate Lane 3: Raji whole cell lysate Lane 4: Daudi whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 57 kDa Blocking/Dilution buffer: 5% NFDM/TBST.





All lanes : Anti-FCRL4 Antibody (C-term) at 1:2000 dilution Lane 1: Daudi whole cell lysate Lane 2: Raji whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 57 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



FCRL4 Antibody (C-term) (Cat. #AP9847b) flow cytometric analysis of Ramos cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

FCRL4 Antibody (C-term) - Background

This gene encodes a member of the immunoglobulin receptor superfamily and is one of several Fc receptor-like glycoproteins clustered on the long arm of chromosome 1. The encoded protein has four extracellular C2-type immunoglobulin domains, a transmembrane domain and a cytoplasmic domain that contains three immune-receptor tyrosine-based inhibitory motifs. This protein may play a role in the function of memory B-cells in the epithelia. Aberrations in the chromosomal region encoding this gene are associated with non-Hodgkin lymphoma and multiple myeloma.

FCRL4 Antibody (C-term) - References

Davila, S., et al. Genes Immun. (2010) In press: Weiss, G.E., et al. J. Immunol. 183(3):2176-2182(2009) Kazemi, T., et al. Cancer Immunol. Immunother. 58(6):989-996(2009) Falini, B., et al. Blood 102(10):3684-3692(2003)