

Anti-IL-13 Reference Antibody (GSK 679586)

Recombinant Antibody Catalog # APR10550

Specification

Anti-IL-13 Reference Antibody (GSK 679586) - Product Information

Application Primary Accession Reactivity Clonality Isotype Calculated MW FC, Kinetics, Animal Model P35225 Human Monoclonal IgG1 145 KDa

Anti-IL-13 Reference Antibody (GSK 679586) - Additional Information

Target/Specificity IL-13

Endotoxin < 0.001EU/ μg,determined by LAL method.

Conjugation Unconjugated

Expression system CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Anti-IL-13 Reference Antibody (GSK 679586) - Protein Information

Name IL13

Synonyms NC30

Function

Cytokine that plays important roles in allergic inflammation and immune response to parasite infection (PubMed:8096327, PubMed:8096327). Synergizes with IL2 in regulating interferon-gamma synthesis (PubMed:8096327). Synergizes with IL2 in regulating interferon-gamma synthesis (PubMed:8096327). Stimulates B-cell proliferation, and activation of eosinophils, basophils, and mast cells (PubMed:7903680, PubMed:8759755). Plays an important role in controlling IL33 activity by modulating the production of transmembrane and soluble forms of interleukin-1 receptor-like 1/IL1RL1 (By similarity). Displays the capacity to



antagonize Th1-driven proinflammatory immune response and downregulates synthesis of many proinflammatory cytokines including IL1, IL6, IL10, IL12 and TNF-alpha through a mechanism that partially involves suppression of NF-kappa-B (By similarity). Also functions on nonhematopoietic cells, including endothelial cells where it induces vascular cell adhesion protein 1/VCAM1, which is important in the recruitment of eosinophils (PubMed:8639787). Exerts its biological effects through its receptors which comprises the IL4R chain and the IL13RA1 chain, to activate JAK1 and TYK2, leading to the activation of STAT6 (PubMed:9013879). Aside from

IL13RA1, another receptor IL13RA2 acts as a high affinity decoy for IL13 and mediates internalization and depletion of extracellular IL13 (PubMed:21622864).


Cellular Location Secreted.

Anti-IL-13 Reference Antibody (GSK 679586) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-IL-13 Reference Antibody (GSK 679586) - Images



Anti-IL-13 Reference Antibody (GSK 679586) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%





The purity of Anti-IL-13 Reference Antibody (GSK 679586) is more than 95% , determined by SEC-HPLC.