

Anti-IL-10 Reference Antibody (BT-063)
Recombinant Antibody
Catalog # APR10630**Specification**

Anti-IL-10 Reference Antibody (BT-063) - Product Information

Application	FC, Kinetics, Animal Model
Primary Accession	P22301
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	145.74 KDa

Anti-IL-10 Reference Antibody (BT-063) - Additional Information**Target/Specificity**
IL-10**Endotoxin**
< 0.001EU/ µg,determined by LAL method.**Conjugation**
Unconjugated**Expression system**
CHO Cell**Format**
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.**Anti-IL-10 Reference Antibody (BT-063) - Protein Information****Name** IL10**Function**
Major immune regulatory cytokine that acts on many cells of the immune system where it has profound anti-inflammatory functions, limiting excessive tissue disruption caused by inflammation. Mechanistically, IL10 binds to its heterotetrameric receptor comprising IL10RA and IL10RB leading to JAK1 and STAT2-mediated phosphorylation of STAT3 (PubMed:16982608). In turn, STAT3 translocates to the nucleus where it drives expression of anti-inflammatory mediators (PubMed:18025162). Targets antigen-presenting cells (APCs) such as macrophages and monocytes and inhibits their release of pro- inflammatory cytokines including granulocyte-macrophage colony- stimulating factor /GM-CSF, granulocyte colony-stimulating factor/G- CSF, IL-1 alpha, IL-1 beta, IL-6, IL-8 and TNF-alpha (PubMed:11564774, PubMed:1940799).

target="_blank">1940799, PubMed:7512027). Also interferes with antigen presentation by reducing the expression of MHC-class II and co- stimulatory molecules, thereby inhibiting their ability to induce T cell activation (PubMed:8144879). In addition, controls the inflammatory response of macrophages by reprogramming essential metabolic pathways including mTOR signaling (By similarity).

Cellular Location
Secreted.


Tissue Location
Produced by a variety of cell lines, including T- cells, macrophages, mast cells and other cell types

Anti-IL-10 Reference Antibody (BT-063) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IL-10 Reference Antibody (BT-063) - Images



Anti-IL-10 Reference Antibody (BT-063) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-IL-10 Reference Antibody (BT-063) is more than 100% ,determined by

SEC-HPLC.