

## Anti-INHBA / Activin A Reference Antibody (garetosmab)

Recombinant Antibody Catalog # APR10679

#### **Specification**

## Anti-INHBA / Activin A Reference Antibody (garetosmab) - Product Information

Application
Primary Accession
Reactivity
Clonality

Isotype Calculated MW FC, Kinetics, Animal Model

P08476

Human, Mouse Monoclonal IgG4SP 145.52 KDa

# Anti-INHBA / Activin A Reference Antibody (garetosmab) - Additional Information

Target/Specificity
INHBA / Activin A

**Endotoxin** 

< 0.001EU/ µg,determined by LAL method.

**Conjugation** Unconjugated

**Expression system** 

CHO Cell

#### **Format**

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

## Anti-INHBA / Activin A Reference Antibody (garetosmab) - Protein Information

#### Name INHBA

#### **Function**

Inhibins/activins are involved in regulating a number of diverse functions such as hypothalamic and pituitary hormone secretion, gonadal hormone secretion, germ cell development and maturation, erythroid differentiation, insulin secretion, nerve cell survival, embryonic axial development or bone growth, depending on their subunit composition.

**Cellular Location** 

Secreted.

#### Anti-INHBA / Activin A Reference Antibody (garetosmab) - Protocols




101. 000.070.1000 Tax. 000.070.100


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## Anti-INHBA / Activin A Reference Antibody (garetosmab) - Images



Anti-INHBA / Activin A Reference Antibody (garetosmab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-INHBA / Activin A Reference Antibody (garetosmab)is more than 91.47% ,determined by SEC-HPLC.