

**Anti-CD74 Reference Antibody (milatuzumab)
Recombinant Antibody
Catalog # APR10818****Specification****Anti-CD74 Reference Antibody (milatuzumab) - Product Information**

Application	FC, Kinetics, Animal Model
Primary Accession	P04233
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	145.66 KDa

Anti-CD74 Reference Antibody (milatuzumab) - Additional Information**Target/Specificity**
CD74**Endotoxin**
< 0.001EU/ µg,determined by LAL method.**Conjugation**
Unconjugated**Expression system**
CHO Cell**Format**
Purified monoclonal antibody supplied in PBS, pH6.0, without preservative.This antibody is purified through a protein A column.**Anti-CD74 Reference Antibody (milatuzumab) - Protein Information****Name** CD74 ([HGNC:1697](#))**Synonyms** DHLAG**Function**
Plays a critical role in MHC class II antigen processing by stabilizing peptide-free class II alpha/beta heterodimers in a complex soon after their synthesis and directing transport of the complex from the endoplasmic reticulum to the endosomal/lysosomal system where the antigen processing and binding of antigenic peptides to MHC class II takes place. Serves as cell surface receptor for the cytokine MIF. [Isoform p41]: Stabilizes the conformation of mature CTSL by binding to its active site and serving as a chaperone to help maintain a pool of mature enzyme in endocytic compartments and extracellular space of antigen-presenting cells (APCs). Has antiviral activity by stymieing the endosomal entry of Ebola virus and coronaviruses, including SARS-CoV-2 (PubMed:32855215). Disrupts cathepsin-mediated Ebola virus glycoprotein processing, which prevents viral fusion and

entry. This antiviral activity is specific to p41 isoform (PubMed:32855215).

Cellular Location

Cell membrane; Single-pass type II membrane protein. Endoplasmic reticulum membrane. Golgi apparatus, trans-Golgi network. Endosome. Lysosome. Secreted. Note=Transits through a number of intracellular compartments in the endocytic pathway. It can either undergo proteolysis or reach the cell membrane

Tissue Location


Detected in urine (at protein level). [Isoform p33]: In B cells, represents 70% of total CD74 expression.

Anti-CD74 Reference Antibody (milatuzumab) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CD74 Reference Antibody (milatuzumab) - Images



Anti-CD74 Reference Antibody (milatuzumab) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%



The purity of Anti-CD74 Reference Antibody (milatuzumab) is more than 100% ,determined by SEC-HPLC.