

CX3CR1 Antibody
Catalog # ASC10081**Specification**

CX3CR1 Antibody - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	P49238
Other Accession	NP_001328 , 4503171
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 39, 43 kDa

Application Notes	Observed: 50 kDa KDa CX3CR1 antibody can be used for detection of CX3CR1 by Western blot at 0.5 µg/mL. Antibody can also be used for immunohistochemistry starting at 10 µg/mL. For immunofluorescence start at 20 µg/mL.
-------------------	--

CX3CR1 Antibody - Additional InformationGene ID **1524****Other Names**

CX3CR1 Antibody: V28, CCRL1, GPR13, CMKDR1, GPRV28, CMKBRL1, CX3C chemokine receptor 1, Beta chemokine receptor-like 1, C-X3-C CKR-1, chemokine (C-X3-C motif) receptor 1

Target/Specificity

CX3CR1; At least two isoforms of CX3CR1 are known to exist.

Reconstitution & Storage

Antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.

Precautions

CX3CR1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CX3CR1 Antibody - Protein Information**Name** CX3CR1 {ECO:0000303|PubMed:12551893, ECO:0000312|HGNC:HGNC:2558}**Function**

Receptor for the C-X3-C chemokine fractalkine (CX3CL1) present on many early leukocyte cells; CX3CR1-CX3CL1 signaling exerts distinct functions in different tissue compartments, such as immune response, inflammation, cell adhesion and chemotaxis (PubMed:12055230, PubMed:12055230).

[23125415](http://www.uniprot.org/citations/23125415), PubMed:<[9390561](http://www.uniprot.org/citations/9390561)>, PubMed:<[9782118](http://www.uniprot.org/citations/9782118)>). CX3CR1-CX3CL1 signaling mediates cell migratory functions (By similarity). Responsible for the recruitment of natural killer (NK) cells to inflamed tissues (By similarity). Acts as a regulator of inflammation process leading to atherogenesis by mediating macrophage and monocyte recruitment to inflamed atherosclerotic plaques, promoting cell survival (By similarity). Involved in airway inflammation by promoting interleukin 2-producing T helper (Th2) cell survival in inflamed lung (By similarity). Involved in the migration of circulating monocytes to non-inflamed tissues, where they differentiate into macrophages and dendritic cells (By similarity). Acts as a negative regulator of angiogenesis, probably by promoting macrophage chemotaxis (PubMed:<[14581400](http://www.uniprot.org/citations/14581400)>, PubMed:<[18971423](http://www.uniprot.org/citations/18971423)>). Plays a key role in brain microglia by regulating inflammatory response in the central nervous system (CNS) and regulating synapse maturation (By similarity). Required to restrain the microglial inflammatory response in the CNS and the resulting parenchymal damage in response to pathological stimuli (By similarity). Involved in brain development by participating in synaptic pruning, a natural process during which brain microglia eliminates extra synapses during postnatal development (By similarity). Synaptic pruning by microglia is required to promote the maturation of circuit connectivity during brain development (By similarity). Acts as an important regulator of the gut microbiota by controlling immunity to intestinal bacteria and fungi (By similarity). Expressed in lamina propria dendritic cells in the small intestine, which form transepithelial dendrites capable of taking up bacteria in order to provide defense against pathogenic bacteria (By similarity). Required to initiate innate and adaptive immune responses against dissemination of commensal fungi (mycobiota) component of the gut: expressed in mononuclear phagocytes (MNPs) and acts by promoting induction of antifungal IgG antibodies response to confer protection against disseminated C.albicans or C.auris infection (PubMed:<[29326275](http://www.uniprot.org/citations/29326275)>). Also acts as a receptor for C-C motif chemokine CCL26, inducing cell chemotaxis (PubMed:<[20974991](http://www.uniprot.org/citations/20974991)>).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location


Expressed in lymphoid and neural tissues (PubMed:7590284). Expressed in lymphocyte subsets, such as natural killer (NK) cells, gamma-delta T-cells and terminally differentiated CD8(+) T-cells (PubMed:12055230). Expressed in smooth muscle cells in atherosclerotic plaques (PubMed:14581400)


CX3CR1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CX3CR1 Antibody - Images






Western blot analysis of CX3CR1 in THP-1 cell lysate with CX3CR1 antibody at 1 μ g/mL in (A) the absence and (B) the presence of blocking peptide.



Immunohistochemistry of CX3CR1 in human spleen cells with CX3CR1 antibody at 10 μ g/mL.



Immunofluorescence of CX3CR1 in THP-1 cells with CX3CR1 antibody at 20 μ g/mL.

CX3CR1 Antibody - Background

CX3CR1 Antibody: CX3CR1 is one of the chemokine receptors that are required as coreceptors for HIV infection. The genes encoding human, mouse, and rat CX3CR1 were cloned and designated V28 and CMKBRL1, CX3CR1, and RBS11, respectively. The encoded seven transmembrane protein was initially identified as the receptor for the transmembrane molecule fractalkine, and renamed CX3CR1. CX3CR1 is known to serve as a coreceptor for HIV-1 and HIV-2 envelope fusion and virus infection, which can be inhibited by fractokine. CX3CR1 mediates leukocyte migration and adhesion. CX3CR1 is expressed in a variety of human tissues and cell lines.

CX3CR1 Antibody - References

Raport CJ, Schweickart VL, Eddy RL JR, et al. The orphan G-protein-coupled receptor-encoding gene V28 is closely related to genes for chemokine receptors and is expressed in lymphoid and neural tissues. *Gene* 1995; 163:295-9

Combadiere C, Ahuja SK, and Murphy PM. Cloning, chromosomal localization, and RNA expression

of a human β chemokine receptor-like gene. DNA Cell Biol. 1995;14:673-80

Combadiere C, Gao J, Tiffany HL, et al. Gene cloning, RNA distribution, and functional expression of mCX3CR1, a mouse chemotactic receptor for the CX3C chemokine fractalkine. Biochem. Biophys. Res. Commun. 1998; 253:728-32

Harrison JK, Barber CM, and Lynch KR. cDNA cloning of a G-protein-coupled receptor expressed in rat spinal cord and brain related to chemokine receptors. Neurosci. Lett. 1994; 169:85-9