

Bcl-rambo Antibody
Catalog # ASC10167**Specification**

Bcl-rambo Antibody - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	Q9B XK5
Other Accession	AAH07658 , 14043326
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	Bcl-rambo antibody can be used for the detection of Bcl-rambo by Western blot at 2 and 4 µg/mL. Antibody can also be used for immunohistochemistry starting at 10 µg/mL. For immunofluorescence start at 20 µg/mL.

Bcl-rambo Antibody - Additional InformationGene ID **23786****Other Names**

Bcl-rambo Antibody: MIL1, BCL-RAMBO, Bcl2-L-13, MIL1, CD003, Bcl-2-like protein 13, Bcl-rambo, BCL2-like 13 (apoptosis facilitator)

Target/Specificity

BCL2L13;

Reconstitution & Storage

Bcl-rambo antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

Bcl-rambo Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Bcl-rambo Antibody - Protein Information**Name** BCL2L13**Synonyms** MIL1**Function**

May promote the activation of caspase-3 and apoptosis.

Cellular Location

[Isoform 2]: Mitochondrion membrane; Single-pass membrane protein. Nucleus

Tissue Location


Ubiquitous, with the highest levels of expression in heart, placenta and pancreas

Bcl-rambo Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Bcl-rambo Antibody - Images



Western blot analysis of Bcl-rambo in K562 cell lysate with Bcl-rambo antibody at (A) 2 and (B) 4 $\mu\text{g/mL}$.



Immunohistochemistry of Bcl-rambo in human heart tissue with Bcl-rambo antibody at 10 $\mu\text{g/mL}$.



Immunofluorescence of Bcl-rambo in Human Heart cells with Bcl-rambo antibody at 20 µg/mL.

Bcl-rambo Antibody - Background

Bcl-rambo Antibody: Apoptosis plays a major role in normal organism development, tissue homeostasis, and removal of damaged cells. Disruption of this process has been implicated in a variety of diseases such as cancer. Members of the Bcl-2 family are known to be critical regulators of this process. These proteins are characterized by the presence of several conserved motifs termed Bcl-2 homology (BH) domains. A novel, widely expressed member termed Bcl-rambo was recently identified. This protein is localized to mitochondria in mammalian cells and its overexpression induces apoptosis which could be blocked by co-expression of inhibitor of apoptosis proteins (IAPs) such as XIAP, cIAP1, and cIAP2. Bcl-rambo shows overall homology to the anti-apoptotic members containing BH motifs, but unlike Bcl-2, the C-terminal membrane anchor of Bcl-rambo is preceded by a unique 250 amino acid insertion. This region by itself can induce apoptosis more efficiently than the Bcl-2 homology regions, suggesting that Bcl-rambo may be important other pro-apoptotic pathways.

Bcl-rambo Antibody - References

- Lockshin RA, Osborne B, and Zakeri Z. Cell death in the third millennium. *Cell Death Differ.* 2000; 7:2-7.
- Cory S, Huang DCS, and Adams JM. The Bcl-2 family: roles in cell survival and oncogenesis. *Oncogene* 2003; 22:8590-607.
- Heiser D, Labi V, Erlacher M, et al. The Bcl-2 protein family and its role in the development of neoplastic disease. *Exp. Gerontol.* 2004; 39:1125-35.
- Kataoka T, Holler N, Michau O, et al. Bcl-rambo, a novel Bcl-2 homologue that induces apoptosis via its unique C-terminal extension. *J. Biol. Chem.* 2001; 276:19548-54.,