

**TAB2 Antibody**  
**Catalog # ASC10267**

## Specification

## TAB2 Antibody - Product Information

Application	IHC-P, E
Primary Accession	<a href="#">Q9NYJ8</a>
Other Accession	<a href="#">NP_055908</a> , <a href="#">14149669</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	TAB2 antibody can be used for detection of TAB2 by immunohistochemistry at 5 µg/mL.

## TAB2 Antibody - Additional Information

## Gene ID

23118

## Other Names

TAB2 Antibody: CHTD2, MAP3K7IP2, KIAA0733, TGF-beta-activated kinase 1 and MAP3K7-binding protein 2, Mitogen-activated protein kinase kinase kinase 7-interacting protein 2, TAB-2, TGF-beta activated kinase 1/MAP3K7 binding protein 2

## Target/Specificity

TAB2; TAB2 antibody is human specific. TAB2 antibody is predicted not to cross-react with other TAB proteins.

## **Reconstitution & Storage**

TAB2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

## Precautions

TAB2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## TAB2 Antibody - Protein Information

**Name TAB2 {ECO:0000303|PubMed:10882101, ECO:0000312|HGNC:HGNC:17075}**

## Function

href="http://www.uniprot.org/citations/33184450" target="\_blank">>33184450</a>, PubMed:<a href="http://www.uniprot.org/citations/36681779" target="\_blank">>36681779</a>). Acts as an adapter linking MAP3K7/TAK1 and TRAF6 to 'Lys-63'-linked polyubiquitin chains (PubMed:<a href="http://www.uniprot.org/citations/10882101" target="\_blank">>10882101</a>, PubMed:<a href="http://www.uniprot.org/citations/11460167" target="\_blank">>11460167</a>, PubMed:<a href="http://www.uniprot.org/citations/15327770" target="\_blank">>15327770</a>, PubMed:<a href="http://www.uniprot.org/citations/22158122" target="\_blank">>22158122</a>, PubMed:<a href="http://www.uniprot.org/citations/27746020" target="\_blank">>27746020</a>). The RanBP2-type zinc finger (NZF) specifically recognizes Lys-63'-linked polyubiquitin chains unanchored or anchored to the substrate proteins such as RIPK1/RIP1 and RIPK2: this acts as a scaffold to organize a large signaling complex to promote autophosphorylation of MAP3K7/TAK1, and subsequent activation of I- kappa-B-kinase (IKK) core complex by MAP3K7/TAK1 (PubMed:<a href="http://www.uniprot.org/citations/15327770" target="\_blank">>15327770</a>, PubMed:<a href="http://www.uniprot.org/citations/18079694" target="\_blank">>18079694</a>, PubMed:<a href="http://www.uniprot.org/citations/22158122" target="\_blank">>22158122</a>). Also recognizes and binds Lys-63'- linked polyubiquitin chains of heterotypic 'Lys-63'-'Lys-48'-linked branched ubiquitin chains (PubMed:<a href="http://www.uniprot.org/citations/27746020" target="\_blank">>27746020</a>). Regulates the IL1-mediated translocation of NCOR1 out of the nucleus (By similarity). Involved in heart development (PubMed:<a href="http://www.uniprot.org/citations/20493459" target="\_blank">>20493459</a>).

### Cellular Location

Membrane; Peripheral membrane protein. Endosome membrane; Peripheral membrane protein. Lysosome membrane; Peripheral membrane protein. Cytoplasm, cytosol. Note=Following IL1 stimulation, translocation occurs from the membrane to cytosol (PubMed:10882101) Interaction with TRIM38 promotes translocation from cytosol to endosome and lysosome (PubMed:24434549).

### Tissue Location


Widely expressed. In the embryo, expressed in the ventricular trabeculae, endothelial cells of the conotruncal cushions of the outflow tract and in the endothelial cells lining the developing aortic valves.

### TAB2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### TAB2 Antibody - Images



Immunohistochemistry of TAB2 in human brain tissue with TAB2 antibody at 5 µg/mL.

#### TAB2 Antibody - Background

**TAB2 Antibody:** TAB2 is an activator of MAP3K7/TAK1, which is required for the IL-1 induced activation NF-kappaB and MAPK8/JNK. This protein forms a kinase complex with TRAF6, MAP3K7 and TAB1, thus serves as an adaptor linking MAP3K7 and TRAF6. This protein, TAB1, and MAP3K7 also participate in the signal transduction induced by TNFSF11/RANKL through the activation of the receptor activator of NF-kappaB (TNFRSF11A/RANK), which may regulate the development and function of osteoclasts. Recent experiments have shown that TAB2 and the related protein TAB3 constitutively interact with the autophagy mediator Beclin-1; upon induction of autophagy, these proteins dissociate from Beclin-1 and bind TAK1. Overexpression of TAB2 and TAB3 inhibit autophagy, while their depletion triggers it, suggesting that TAB2 and TAB3 act as a control point for autophagy.

#### TAB2 Antibody - References

- Takaesu G, Kishida S, Hiyama A, et al. TAB2, a novel adaptor protein, mediates activation of TAK1 MAPKKK by linking TAK1 to TRAF6 in the IL-1 signal transduction pathway. *Mol. Cell* 2000; 5:649-58.  
Maizukami J, Takaesu G, Akatsuka H, et al. Receptor activation of NF-kappaB ligand (RANKL) activates TAK1 mitogen-activated protein kinase kinase kinase through a signaling complex containing RANK, TAB2, and TRAF6. *Mol. Cell. Biol.* 2002; 22:992-1000.  
Criollo A, Niso-Santano M, Malik SA, et al. Inhibition of autophagy by TAB2 and TAB3. *EMBO J.* 2011; 30:4908-20.