

FEZ2 Antibody
Catalog # ASC10616**Specification**

FEZ2 Antibody - Product Information

Application	WB, IF, E
Primary Accession	Q9UHY8
Other Accession	NP_001036013 , 110349756
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	FEZ2 antibody can be used for the detection of FEZ2 by Western blot at 0.5 - 1 µg/mL. Antibody can also be used for immunofluorescence starting at 5 µg/mL. For immunofluorescence start at 5 µg/mL.

FEZ2 Antibody - Additional Information

Gene ID	9637
Target/Specificity	
FEZ2;	

Reconstitution & Storage

FEZ2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

FEZ2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

FEZ2 Antibody - Protein Information

Name FEZ2

Function

Involved in axonal outgrowth and fasciculation.

Tissue Location


Expressed in nonneural tissues, such as heart, lung, spleen, muscle, testis, placenta and melanocytes

FEZ2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FEZ2 Antibody - Images



Western blot analysis of FEZ2 in 3T3 cell lysate with FEZ2 antibody at (A) 0.5, and (B) 1 µg/mL.



Immunofluorescence of human brain tissue using FEZ2 antibody at 5 µg/mL.

FEZ2 Antibody - Background

FEZ2 Antibody: Fasciculation and elongation protein zeta-2 (FEZ2) is a homolog to the mammalian FEZ1, itself an ortholog of the *C. elegans* UNC-76. In contrast to FEZ1, FEZ2 mRNA is widely expressed in mouse tissues. FEZ2 interacts with protein kinase C (PKC)-zeta substrate and induces neurite extension of PC12 cells when co-expressed with a constitutively active form of PKC-zeta, suggesting FEZ2 may play an important role in the morphological changes of various cells by associating with PKC-zeta in a tissue non-specific manner. FEZ2 can interact with FEZ1 through its c-terminal regions and especially its coiled-coil region. At least two isoforms of FEZ2 are known to exist.

FEZ2 Antibody - References

Fujita T, Ikuta J, Okajima T, et al. Identification of a tissue-non-specific homologue of axonal fasciculation and elongation protein zeta-1. *Biochem. Biophys. Res. Commun.*2004; 313:738-44.
Kuroda S, Nakagawa N, Tokunaga C, et al. Mammalian homologue of the *Caenorhabditis elegans*

UNC-76 protein involved in axonal outgrowth is a protein kinase C ζ -interacting protein. J. Cell Biol.1999; 144:403-11.

Assmann EM, Alborghetti MR, Camargo MER, et al. FEZ1 dimerization and interaction with transcription regulatory proteins involves its coiled-coil region. J. Biol. Chem.2006; 281:9869-81.