

MC4R Antibody
Catalog # ASC10917**Specification**

MC4R Antibody - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	P32245
Other Accession	NP_005903 , 119508433
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	MC4R antibody can be used for detection of MC4R by Western blot at 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 20 µg/mL.

MC4R Antibody - Additional Information

Gene ID	4160
Target/Specificity	
MC4R;	

Reconstitution & Storage

MC4R antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

MC4R Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

MC4R Antibody - Protein Information

Name MC4R

Function

Hormone receptor that acts as a key component of the leptin- melanocortin pathway at the intersection of homeostatic maintenance of energetic state (PubMed:32327598, PubMed:33858992). Plays a role in regulating food intake: activation by a stimulating hormone such as anorexigenic alpha-melanocyte stimulating hormone (alpha-MSH) inhibits appetite, whereas binding to a natural antagonist like Agouti-related protein/AGRP promotes appetite. G-protein-coupled receptor that activates conventional Gα_s signaling leading to induction of anorexigenic signaling in the hypothalamus to result in negative energy balance (PubMed:33858992). Regulates

the firing activity of neurons from the hypothalamus by alpha-MSH and AGRP independently of Galphas signaling by ligand-induced coupling of closure of inwardly rectifying potassium channel KCNJ13 (By similarity). In intestinal epithelial cells, plays a role in the inhibition of hepatic glucose production via nesfatin-1/NUCB2 leading to increased cyclic adenosine monophosphate (cAMP) levels and glucagon-like peptide 1 (GLP-1) secretion in the intestinal epithelium (PubMed:39562740).

Cellular Location

Cell membrane; Multi-pass membrane protein


Tissue Location

Brain, placental, and gut tissues.


MC4R Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MC4R Antibody - Images

Western blot analysis of MC4R in rat brain tissue lysate with MC4R antibody at 1 µg/mL in (A) the absence and (B) the presence of blocking peptide.



Immunohistochemistry of MC4R in human brain tissue with MC4R antibody at 2.5 µg/mL.



Immunofluorescence of MC4R in Human Brain cells with MC4R antibody at 20 µg/mL.

MC4R Antibody - Background

MC4R Antibody: The melanocortin-4 receptor (MC4R) is a member of the superfamily of seven transmembrane G-protein coupled receptors that are involved in multiple signal transduction pathways including the cAMP and MAPK signaling pathways. It is thought that the melanocortin system modulates energy expenditure and insulin sensitivity; activation of the MC4R results in the inhibition of c-Jun N-terminal kinase (JNK) activity and promotes insulin signaling. MC4R-null mice display maturity onset obesity characterized by hyperphagia, increased adiposity, hyperinsulinaemia and hyperleptinaemia, suggesting that like other obesity-linked genes such as FTO, PTER, and NPC1, MC4R is a potential candidate target for the treatment of obesity.

MC4R Antibody - References

Gantz I, Miwa H, Konda Y, et al. Molecular cloning, expression, and gene localization of a fourth melanocortin receptor. *J. Biol. Chem.*1993; 268:15174-9.
Vongs A, Lynn NM, and Rosenblum CI. Activation of MAP kinase by MC4-R through PI3 kinase. *Regul. Pept.*2004; 120:113-8.
Cone H. Anatomy and regulation of the central melanocortin system. *Nat. Neurosci.*2005; 8:571-8.
Chai B, Li J-Y, Zhang W, et al. Melanocortin-4 receptor activation inhibits c-Jun N-terminal kinase activity and promotes insulin signaling. *Peptides*2009; 30:1098-10