

**SRPX2 Antibody**  
**Catalog # ASC11012****Specification**

---

**SRPX2 Antibody - Product Information**

Application	WB, IHC-P, IF, E
Primary Accession	<a href="#">O60687</a>
Other Accession	<a href="#">NP_055282</a> , <a href="#">7657619</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	SRPX2 antibody can be used for detection of SRPX2 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.

**SRPX2 Antibody - Additional Information**

Gene ID	27286
Target/Specificity	
SRPX2;	

**Reconstitution & Storage**

SRPX2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Precautions**

SRPX2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**SRPX2 Antibody - Protein Information**

**Name** SRPX2

**Synonyms** SRPUL

**Function**

Acts as a ligand for the urokinase plasminogen activator surface receptor. Plays a role in angiogenesis by inducing endothelial cell migration and the formation of vascular network (cords). Involved in cellular migration and adhesion. Increases the phosphorylation levels of FAK. Interacts with and increases the mitogenic activity of HGF. Promotes synapse formation. May have a role in the perisylvian region, critical for language and cognitive development.

**Cellular Location**

Secreted. Cytoplasm. Cell surface. Synapse

### Tissue Location


Expressed in neurons of the rolandic area of the brain (at protein level). Highly expressed in the brain, placenta, lung, trachea, uterus, adrenal gland, heart, ovary and placenta. Weakly expressed in the peripheral blood, brain and bone marrow. Expressed in numerous cancer cell lines and in gastrointestinal cancer cells. Higher levels found in colorectal cancers than in normal colonic mucosa

### SRPX2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### SRPX2 Antibody - Images



Western blot analysis of SRPX2 in human lung tissue lysate with SRPX2 antibody at (A) 1 and (B) 2  $\mu$ g/mL.



Immunohistochemistry of SRPX2 in rat lung tissue with SRPX2 antibody at 5 µg/mL.



Immunofluorescence of SRPX2 in human lung tissue with SRPX2 antibody at 20 µg/mL.

### **SRPX2 Antibody - Background**

**SRPX2 Antibody:** Sushi-repeat-containing protein X-linked 2 (SRPX2) is a neural gene functioning in the speech and language center of the human brain; mutations in this gene lead to epilepsy, speech dyspraxia, mental retardation and cognitive disorders. Recently, SRPX2 was found to be a novel mediator of angiogenesis and can act as a ligand for the urokinase-type plasminogen activator, a protein that can facilitate invasive migration of sprouting endothelial cells. SRPX2 is also overexpressed in gastric cancer, leading to increased phosphorylation levels of focal adhesion kinase and enhanced cellular migration and adhesion, suggesting that SRPX2 may be a potential target in the treatment of metastatic cancers.

### **SRPX2 Antibody - References**

Roll P, Rudolf G, Pereira S, et al. SRPX2 mutations in disorders of language cortex and cognition. Hum. Mol. Genet.2006; 15:1195-207.  
Miljkovic-Licina M, Hammel P, Garrido-Urbani S, et al. Sushi repeat protein X-linked 2, a novel mediator of angiogenesis. FASEB J.2009; 23:4105-16.  
Blasi F and Carmeliet P. uPAR: a versatile signalling orchestrator. Nat. Rev. Mol. Cell. Biol.2002; 3:932-43.  
Tanaka K, Arao T, Maegawa M, et al. SRPX2 is overexpressed in gastric cancer and promotes cellular migration and adhesion. Int. J. Cancer2009; 124:1072-80.