

PANX1 Antibody
Catalog # ASC11574**Specification****PANX1 Antibody - Product Information**

| | |
|-------------------|---|
| Application | WB, IF, E |
| Primary Accession | Q96RD7 |
| Other Accession | NP_056183 , 39995064 |
| Reactivity | Human, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Calculated MW | 47 kDa KDa |
| Application Notes | PANX1 antibody can be used for detection of PANX1 by Western blot at 1 - 2 µg/mL. For immunofluorescence start at 20 µg/mL. |

PANX1 Antibody - Additional InformationGene ID **24145****Target/Specificity**

PANX1; Two transcript variants encoding different isoforms have been found for this gene.

Reconstitution & Storage

PANX1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

PANX1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PANX1 Antibody - Protein InformationName PANX1 ([HGNC:8599](#))**Function**

Ion channel involved in a variety of physiological functions such as blood pressure regulation, apoptotic cell clearance and oogenesis (PubMed:15304325, PubMed:16908669, PubMed:20829356, PubMed:20944749, PubMed:30918116). Forms anion-selective channels with relatively low conductance and an order of permeabilities: nitrate>iodide>chloride>>aspartate=glutamate=gluconate (By similarity). Can release ATP upon activation through phosphorylation or cleavage at C-terminus (PubMed:32238926). May play a role as a Ca(2+)- leak channel to regulate ER Ca(2+) homeostasis (PubMed:16908669).

Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000255|PROSITE-ProRule:PRU00351}.
Endoplasmic reticulum membrane; Multi-pass membrane protein
{ECO:0000255|PROSITE-ProRule:PRU00351}

Tissue Location


Widely expressed (PubMed:30918116). Highest expression is observed in oocytes and brain (PubMed:30918116). Detected at very low levels in sperm cells (PubMed:30918116)

PANX1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PANX1 Antibody - Images



Western blot analysis of PANX1 in human ovary tissue lysate with PANX1 antibody at 1 µg/mL.



Immunofluorescence of PANX1 in human ovary tissue with PANX1 antibody at 20 µg/mL.

PANX1 Antibody - Background

PANX1 Antibody: The pannexin gene family encodes a second class of putative gap junction proteins and are highly conserved in invertebrates and mammals. Pannexins (Panx) are four-pass transmembrane proteins that oligomerize to form large pore ion and metabolite-permeable channels. Pannexin-1 (PANX1) and Pannexin-3 are closely related, while Pannexin-2 is a more distant relation. PANX1 is a transmembrane protein that forms a mechanosensitive ATP-permeable channel between adjacent cells and in the endoplasmic reticulum. PANX1 may play a role as a Ca^{2+} -leak channel to regulate ER Ca^{2+} homeostasis and regulates neural stem and progenitor cell proliferation.

PANX1 Antibody - References

Barbe MT, Monyer H and Bruzzone R. Cell-cell communication beyond connexins: the pannexin channels. *Physiology* 2006; 21:103-14.
Baranova A, Ivanov D, Petrash N, et al. The mammalian pannexin family is homologous to the invertebrate innexin gap junction proteins. *Genomics* 2004; 83:706-16.
Sohl G, Maxeiner S and Willecke K. Expression and functions of neuronal gap junctions. *Nat. Rev. Neurosci.* 2005; 6:191-200
Bao L, Locovei S and Dahl G. Pannexin membrane channels are mechanosensitive conduits for ATP. *FEBS Lett.* 2004; 572:65-8.