

CCL17 Antibody
Catalog # ASC11619**Specification**

CCL17 Antibody - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	Q92583
Other Accession	NP_002978 , 4506829
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	10 kDa KDa
Application Notes	CCL17 antibody can be used for detection of CCL17 by Western blot at 1 - 2 µg/mL.

CCL17 Antibody - Additional InformationGene ID **6361****Target/Specificity**
CCL17;**Reconstitution & Storage**

CCL17 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions

CCL17 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CCL17 Antibody - Protein Information**Name** CCL17**Synonyms** SCYA17, TARC**Function**

Chemokine, which displays chemotactic activity for T lymphocytes, preferentially Th2 cells, but not monocytes or granulocytes. Therefore plays an important role in a wide range of inflammatory and immunological processes (PubMed:<[a href="http://www.uniprot.org/citations/8702936" target="_blank">8702936a href="http://www.uniprot.org/citations/9169480" target="_blank">9169480a href="http://www.uniprot.org/citations/10540332" target="_blank">10540332a href="http://www.uniprot.org/citations/9169480" target="_blank">9169480a href="http://www.uniprot.org/citations/27525438" target="_blank">27525438](http://www.uniprot.org/citations/8702936)

(By similarity).

Cellular Location
Secreted

Tissue Location


Constitutively expressed in thymus. Detected at lower levels in the lung, colon and small intestine (PubMed:8702936) Expressed in stimulated peripheral blood mononuclear cells, but not in resting cells (PubMed:8702936).

CCL17 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CCL17 Antibody - Images



Western blot analysis of CCL17 in rat spleen tissue lysate with CCL17 antibody at (A) 1 and (B) 2 μ g/mL



Immunohistochemistry of CCL17 in spleen tissue with CCL17 antibody at 5 µg/ml.



Immunofluorescence of CCL17 in human spleen tissue with CCL17 antibody at 20 µg/ml.

CCL17 Antibody - Background

CCL17 Antibody: CCL17, also known as TARC, binds to chemokine receptors CCR4 and CCR8 and plays important roles in T cell development in thymus as well as in trafficking and activation of mature T cells. CCL17 is secreted from monocyte-derived DCs and endothelial cells and is responsible for selective recruitment and migration of activated Th2 lymphocytes. CCL17 is also required for CCR7 and CXCR4-dependent migration of cutaneous dendritic cells under inflammatory conditions.

CCL17 Antibody - References

- Imai T, Baba M, Nishimura M, et al. The T cell-directed CC chemokine TARC is a highly specific biological ligand for CC chemokine receptor 4. *J. Biol. Chem.* 1997; 272:15036-42.
- Bernardini G, Hedrick J, Sozzani S, et al. Identification of the CC chemokines TARC and macrophage inflammatory protein-1 beta as novel functional ligands for the CCR8 receptor. *Eur. J. Immunol.* 1998; 28:582-8.
- Alferink J, Lieberam I, Reindl W, et al. Compartmentalized production of CCL17 in vivo: strong inducibility in peripheral dendritic cells contrasts selective absence from the spleen. *J. Exp. Med.* 2003; 197:585-99.
- Stutte S, Quast T, Gerbitzki N, et al. Requirement of CCL17 for CCR7 and CXCR4-dependent migration of cutaneous dendritic cells. *Proc. Natl. Acad. Sci. USA* 2010; 107:8736-41.