

CRTC2 Antibody
Catalog # ASC11650**Specification**

CRTC2 Antibody - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	Q53ET0
Other Accession	NP_859066 , 32171215
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 67, 76 kDa
Application Notes	Observed: 70 kDa KDa CRTC2 antibody can be used for detection of CRTC2 by Western blot at 1 - 2 µg/mL.

CRTC2 Antibody - Additional InformationGene ID **200186****Target/Specificity**

CRTC2; Multiple isoforms of CRTC2 are known to exist. CRTC2 antibody is predicted to not cross-react with CRTC1.

Reconstitution & Storage

CRTC2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions

CRTC2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CRTC2 Antibody - Protein Information**Name** CRTC2**Synonyms** TORC2**Function**

Transcriptional coactivator for CREB1 which activates transcription through both consensus and variant cAMP response element (CRE) sites. Acts as a coactivator, in the SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of CREB1 'Ser-133' phosphorylation. Enhances the interaction of CREB1 with TAF4. Regulates gluconeogenesis as a component of the LKB1/AMPK/TORC2 signaling pathway. Regulates the expression of specific genes such as the steroidogenic gene, StAR. Potent coactivator of PPARGC1A and inducer of mitochondrial biogenesis in muscle cells. Also coactivator for TAX activation of the human T-cell leukemia virus type 1 (HTLV-1) long terminal repeats (LTR).

Cellular Location

Cytoplasm. Nucleus. Note=Translocated from the nucleus to the cytoplasm on interaction of the phosphorylated form with 14-3-3 protein (PubMed:15454081). In response to cAMP levels and glucagon, relocated to the nucleus (PubMed:15454081)

Tissue Location


Most abundantly expressed in the thymus. Present in both B and T-lymphocytes. Highly expressed in HEK293T cells and in insulinomas. High levels also in spleen, ovary, muscle and lung, with highest levels in muscle. Lower levels found in brain, colon, heart, kidney, prostate, small intestine and stomach. Weak expression in liver and pancreas.

CRTC2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CRTC2 Antibody - Images



Western blot analysis of CRTC2 in human small intestine tissue lysate with CRTC2 antibody at (A) 0.5 and (B) 1 µg/mL.



Immunohistochemistry of CRTC2 in human small intestine tissue with CRTC2 antibody at 2.5 µg/ml.



Immunofluorescence of CRTC2 in human small intestine tissue with CRTC2 antibody at 20 µg/ml.

CRTC2 Antibody - Background

CRTC2 Antibody: CRTC2, also known as TORC2 (Transducers Of Regulated cAMP Response Element-Binding (CREB)) 2 and the related protein CRTC1 are potent CREB coactivators that are exported from the nucleus in a CRM1-dependent manner via phosphorylation-dependent interactions. Studies suggest that their phosphorylation and nuclear/cytoplasmic shuttling play an important role in the regulation of gluconeogenesis by cAMP. CRTC2 is present in both B and T-lymphocytes and abundantly expressed in the thymus. Its activity is important in regulating the expression of genes involved in cellular energy metabolism while CRTC1 is essential for energy balance and fertility.

CRTC2 Antibody - References

- Conkright MD, Canettieri G, Screaton R, et al. TORCs: transducers of regulated CREB activity. *Mol. Cell* 2003; 12:413-23.
- Iourgenko V, Zhang W, Mickanin C, et al. Identification of a family of cAMP response element-binding protein coactivators by genome-scale functional analysis in mammalian cells. *Proc. Natl. Acad. Sci. USA* 2003; 100:12147-52.
- Bittinger MA, McWhinnie E, Meltzer J, et al. Activation of cAMP response element-mediated gene expression by regulated nuclear transport of TORC proteins. *Curr. Biol.* 2004; 14: 2156-61.
- Jiang S, Inada T, Tanaka M, et al. Involvement of TORC2, a CREB co-activator, in the in vivo-specific

transcriptional control of HTLV-1. Retrovirology 2009; 6:73.