

HDAC1 Antibody
Catalog # ASC11824**Specification****HDAC1 Antibody - Product Information**

| | |
|-------------------|---|
| Application | WB, E |
| Primary Accession | Q13547 |
| Other Accession | NP_004955 , 13128860 |
| Reactivity | Human, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Calculated MW | Predicted: 53 kDa |
| Application Notes | Observed: 54 kDa KDa HDAC1 antibody can be used for detection of HDAC1 by Western blot at 1 - 2 µg/ml. |

HDAC1 Antibody - Additional Information

Gene ID 3065

Target/Specificity

HDAC1; HDAC1 antibody is human, mouse and rat reactive. HDAC1 antibody is predicted to not cross-react with other members of the HDAC family.

Reconstitution & Storage

HDAC1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions

HDAC1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

HDAC1 Antibody - Protein Information

Name HDAC1 {ECO:0000303|PubMed:10846170, ECO:0000312|HGNC:HGNC:4852}

Function

Histone deacetylase that catalyzes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4) (PubMed:[16762839](http://www.uniprot.org/citations/16762839), PubMed:[17704056](http://www.uniprot.org/citations/17704056), PubMed:[28497810](http://www.uniprot.org/citations/28497810)). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (PubMed:[16762839](http://www.uniprot.org/citations/16762839), PubMed:[17704056](http://www.uniprot.org/citations/17704056)). Histone deacetylases act via the formation of large multiprotein complexes (PubMed:[16762839](http://www.uniprot.org/citations/16762839), PubMed:[17704056](http://www.uniprot.org/citations/17704056)). Acts as a

component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin (PubMed: [16428440](http://www.uniprot.org/citations/16428440), PubMed: [28977666](http://www.uniprot.org/citations/28977666)). As part of the SIN3B complex is recruited downstream of the constitutively active genes transcriptional start sites through interaction with histones and mitigates histone acetylation and RNA polymerase II progression within transcribed regions contributing to the regulation of transcription (PubMed: [21041482](http://www.uniprot.org/citations/21041482)). Also functions as a deacetylase for non-histone targets, such as NR1D2, RELA, SP1, SP3, STAT3 and TSHZ3 (PubMed: [12837748](http://www.uniprot.org/citations/12837748), PubMed: [16285960](http://www.uniprot.org/citations/16285960), PubMed: [16478997](http://www.uniprot.org/citations/16478997), PubMed: [17996965](http://www.uniprot.org/citations/17996965), PubMed: [19343227](http://www.uniprot.org/citations/19343227)). Deacetylates SP proteins, SP1 and SP3, and regulates their function (PubMed: [12837748](http://www.uniprot.org/citations/12837748), PubMed: [16478997](http://www.uniprot.org/citations/16478997), PubMed: [19081374](http://www.uniprot.org/citations/19081374)). Upon calcium stimulation, HDAC1 is released from the complex and CREBBP is recruited, which facilitates transcriptional activation (PubMed: [19081374](http://www.uniprot.org/citations/19081374)). Deacetylates TSHZ3 and regulates its transcriptional repressor activity (PubMed: [19343227](http://www.uniprot.org/citations/19343227)). Deacetylates 'Lys-310' in RELA and thereby inhibits the transcriptional activity of NF-kappa-B (PubMed: [17000776](http://www.uniprot.org/citations/17000776)). Deacetylates NR1D2 and abrogates the effect of KAT5- mediated relieving of NR1D2 transcription repression activity (PubMed: [17996965](http://www.uniprot.org/citations/17996965)). Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development (By similarity). Involved in CIART-mediated transcriptional repression of the circadian transcriptional activator: CLOCK-BMAL1 heterodimer (By similarity). Required for the transcriptional repression of circadian target genes, such as PER1, mediated by the large PER complex or CRY1 through histone deacetylation (By similarity). In addition to protein deacetylase activity, also has protein-lysine deacylase activity: acts as a protein decrotonylase and delactylase by mediating decrotonylation ((2E)-butenoyl) and delactylation (lactoyl) of histones, respectively (PubMed: [28497810](http://www.uniprot.org/citations/28497810), PubMed: [35044827](http://www.uniprot.org/citations/35044827)).

Cellular Location

Nucleus

Tissue Location

Ubiquitous, with higher levels in heart, pancreas and testis, and lower levels in kidney and brain


HDAC1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HDAC1 Antibody - Images



Western blot analysis of HDAC1 in human brain tissue lysate with HDAC1 antibody at 1 μ g/ml in (A) the absence and (B) the presence of blocking peptide.

HDAC1 Antibody - Background

The histone deacetylase (HDAC) family contains multiple members which are divided into four classes. Class I of the HDAC family comprises four members, HDAC1, 2, 3, and 8, each of which contains a deacetylase domain and exhibits a different, individual substrate specificity and function in vivo (1). HDAC1 is responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4) (1,2). HDAC1 gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (3,4).

HDAC1 Antibody - References

- Taunton J, Hassig CA, and Schreiber SL. A mammalian histone deacetylase related to the yeast transcriptional regulator Rpd3p. *Science* 1996; 272:408-11.
- Cai RL, Yan-Neale Y, Cueto MA, et al. HDAC1, a histone deacetylase, forms a complex with Hus1 and Rad9, two G2/M checkpoint Rad proteins. *J. Biol. Chem.* 2000; 275:27909-16.
- Winter M, Moser MA, Meunier D, et al. Divergent roles of HDAC1 and HDAC2 in the regulation of epidermal development and tumorigenesis. *EMBO J.* 2013; 32:3176-91.
- Turgeon N, Blais M, Gagné JM, et al. HDAC1 and HDAC2 restrain the intestinal inflammatory response by regulating intestinal epithelial cell differentiation. *PLoS One* 2013; 8:e73785.