

UVRAG Antibody
Catalog # ASM10494**Specification****UVRAG Antibody - Product Information**

Application	WB, ICC
Primary Accession	Q9P2Y5
Other Accession	NP_003360.2
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal

Description

Rabbit Anti-Human UVRAG Polyclonal

Target/Specificity

Predicted molecular weight at ~78.1kDa. Observed molecular weights in the 75-90kDa range.

Other Names

DHTX Antibody, p63 Antibody, UV radiation resistance associated Antibody, UVRAG_HUMAN Antibody

Immunogen

Synthetic peptide from the C-terminal of human UVRAG

Purification

Peptide Affinity Purified

Storage **-20°C****Storage Buffer**

PBS, 50% glycerol, 0.09% sodium azide

Shipping Temperature **Blue Ice or 4°C****Certificate of Analysis**

A 1:1000 dilution of SPC-605 was sufficient for detection of UVRAG on 293T Rapamycin treated lysates using Goat anti-rabbit IgG:HRP as the secondary antibody.

Cellular Localization

Endosome | Lysosome | Endoplasmic reticulum | Nucleus | Chromosome | Centromere


UVRAG Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)


UVRAG Antibody - Images



Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-UVRAG Polyclonal Antibody (ASM10494). Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Rabbit Anti-UVRAG Polyclonal Antibody (ASM10494) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Rabbit ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Late Endosome, Lysosome, Early Endosome. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) UVAG Antibody (D) Composite.



Western blot analysis of Human 293T showing detection of ~70kDa UVAG protein using Rabbit Anti-UVAG Polyclonal Antibody (ASM10494). Lane 1: MW Ladder. Lane 2: Human 293T (20 μg). Load: 20 μg. Block: 5% milk + TBST for 1 hour at RT. Primary Antibody: Rabbit Anti-UVAG Polyclonal Antibody (ASM10494) at 1:1000 for 1 hour at RT. Secondary Antibody: Goat Anti-Rabbit: HRP at 1:2000 for 1 hour at RT. Color Development: TMB solution for 12 min at RT. Predicted/Observed Size: ~70kDa.



UVRAG Antibody - Background

UVRAG (UV radiation resistance-associated gene) is associated with the Beclin-1/PI3KC3 complex and promotes PI3KC3 enzymatic activity and autophagy, while suppressing proliferation (1). Beclin-1 binding to UVRAG promotes both autophagosome maturation and endocytic trafficking (2). UVRAG is also a potential tumor suppressor protein with frameshift mutations observed in colon and gastric carcinomas (3-4). It is highly expressed in the brain, lung, kidney and liver.

UVRAG Antibody - References

1. Liang, C. et al. (2008) *Nat Cell Biol.* 10: 776-87.
2. Ionov, Y. et al. (2004) *Oncogene.* 23: 639-45.
3. Kim, M.S. et al. (2008) *Hum Pathol.* 39: 1059-63.