

Fab Anti-Human IgG (H&L) Secondary Antibody
Goat Polyclonal, Unconjugated
Catalog # ASR1247**Specification****Fab Anti-Human IgG (H&L) Secondary Antibody - Product Information**

Description	F(ab) Anti-HUMAN IgG (H&L) (GOAT) Antibody
Host	Goat
Conjugate	Unconjugated
Target Species	Human
Reactivity	Human
Clonality	Polyclonal
Application	WB, E, IC
Application Note	ELISA 1:40,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000
Physical State	Liquid (sterile filtered)
Host Isotype	IgG F(ab)
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Anti-Human IgG whole molecule was produced by repeated immunization with Human IgG whole molecule in goat.
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide

Fab Anti-Human IgG (H&L) Secondary Antibody - Additional Information**Shipping Condition**

Wet Ice

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum. No reaction was observed against anti-Pepsin or anti-Goat IgG F(c).

Storage Condition

Store vial at 4° C prior to opening. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Fab Anti-Human IgG (H&L) Secondary Antibody - Protein Information

Fab Anti-Human IgG (H&L) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Fab Anti-Human IgG (H&L) Secondary Antibody - Images

Fab Anti-Human IgG (H&L) Secondary Antibody - Background

Anti-Human IgG whole molecule antibody generated in goat detects specifically human IgG whole molecule. This secondary antibody anti-Human is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays.