

Goat IgG (Agarose Conjugated)

Catalog # ASR1268

Specification

Goat IgG (Agarose Conjugated) - Product Information

Description GOAT IgG whole molecule Agarose

Conjugated Unconjugated

Physical State Suspension of agarose beads

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Species of Origin
Stabilizer

Goat
None

Preservative 0.05% (w/v) Sodium Azide

Goat IgG (Agarose Conjugated) - Additional Information

Shipping Condition

Wet Ice

Purity

This product is normal Goat IgG coupled to activated agarose. A single precipitin arc was observed against anti-Goat Serum when assayed by immunoelectrophoresis prior to coupling to the beads.

Storage Condition

Store vial at 4° C prior to opening. DO NOT FREEZE.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Goat IgG (Agarose Conjugated) - Protein Information


Goat IgG (Agarose Conjugated) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Goat IgG (Agarose Conjugated) - Images





SDS-PAGE of Goat IgG Whole Molecule Agarose Conjugated . Lane M: 5 μ L Opal Prestained Marker . Lane 1: Reduced Goat IgG Whole Molecule Agarose Conjugated . Lane 2: Reduced Goat IgG F(c) Fragment . Lane 3: Reduced Goat IgG F(ab) Fragment . Lane 4: Reduced Goat IgM Whole Molecule . Load: 1 μ g for IgG, F(c) and F(ab); 3 μ g for IgM. Predicted/Observed size: IgG at 50 and 25 kDa; F(c) at 25 kDa; F(ab) at 25 kDa; IgM at 70 and 23 kDa. Observed F(c) Fragment migrates slightly higher.