

# Anti-Goat IgG (H&L) Secondary Antibody

Mouse Polyclonal, Unconjugated Catalog # ASR1325

### Specification

# Anti-Goat IgG (H&L) Secondary Antibody - Product Information

Description Host Conjugate Target Species Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen Reconstitution Volume Reconstitution Buffer

Stabilizer Preservative Anti-GOAT IgG (H&L) (MOUSE) Antibody Mouse Unconjugated Goat Polyclonal WB, E, IC ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000 Lvophilized Antiserum IaG (H&L) 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Goat IgG whole molecule 2.0 mL Restore with deionized water (or equivalent) None None

### Anti-Goat IgG (H&L) Secondary Antibody - Additional Information

Shipping Condition Ambient

**Purity** 

This product was prepared from monospecific ascites by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against Goat IgG and Goat Serum.

#### Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note** This product is for research use only and is not intended for therapeutic or diagnostic applications.

### Anti-Goat IgG (H&L) Secondary Antibody - Protein Information



# Anti-Goat IgG (H&L) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Goat IgG (H&L) Secondary Antibody - Images