

# Anti-Goat IgM (mu) Secondary Antibody

Rabbit Polyclonal, Unconjugated Catalog # ASR1329

### **Specification**

**Physical State** 

**Immunogen** 

### Anti-Goat IgM (mu) Secondary Antibody - Product Information

Description Anti-GOAT IgM (mu) (RABBIT) Antibody

Host Rabbit

Conjugate Unconjugated

Target Species
Clonality
Application

Goat
Polyclonal
WB, E, IC

Application Note ELISA 1:20,000-1:100,000; Western Blot

1:2,000-1:10,000;Immunochemistry

1:1,000-1:5,000 Lyophilized Antiserum

 $\begin{array}{ccc} \text{Host Isotype} & & \text{Antiserum} \\ \text{Target Isotype} & & \text{IgM } \mu \text{ chain} \end{array}$ 

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2
Goat IgM mu heavy chain

Reconstitution Volume 2.0 m

Reconstitution Buffer Restore with deionized water (or

equivalent)

# Anti-Goat IgM (mu) Secondary Antibody - Additional Information

## **Shipping Condition**

**Ambient** 

#### **Purity**

This product was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against Goat IgM and Goat Serum. No reaction was observed against Goat IgG.

#### **Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

# Anti-Goat IgM (mu) Secondary Antibody - Protein Information



Tel: 858.875.1900 Fax: 858.875.1999



# Anti-Goat IgM (mu) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Goat IgM (mu) Secondary Antibody - Images