

Anti-Guinea Pig IgG F(c) Secondary Antibody

Goat Polyclonal, Unconjugated Catalog # ASR1392

Specification

Anti-Guinea Pig IgG F(c) Secondary Antibody - Product Information

Description Host Conjugate Target Species Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen Stabilizer Preservative Anti-GUINEA PIG IgG F(c) (GOAT) Antibody Goat Unconjugated **Guinea** Pig Polyclonal WB, E, IC ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000 Liquid (sterile filtered) laG laG F(c) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Guinea Pig IgG F(c) fragment None 0.01% (w/v) Sodium Azide

Anti-Guinea Pig IgG F(c) Secondary Antibody - Additional Information

Shipping Condition Wet Ice

Purity

Anti-Guinea Pig IgG F(c) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Guinea Pig IgG, Guinea Pig IgG F(c) and Guinea Pig Serum. No reaction was observed against Guinea Pig IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Guinea Pig IgG F(c) Secondary Antibody - Protein Information



Anti-Guinea Pig IgG F(c) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Guinea Pig IgG F(c) Secondary Antibody - Images

Anti-Guinea Pig IgG F(c) Secondary Antibody - Background

Anti-Guinea Pig IgG F(c) antibody generated in Goat detects specifically Guinea Pig IgG F(c). Anti-Guinea Pig IgG F(c) secondary antibody is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays.Anti-Guinea Pig IgG F(c) antibody is ideal for investigators in Cancer, Immunology, and Microbiology research.