

Anti-Horse IgG F(c) Secondary Antibody
Goat Polyclonal, Unconjugated
Catalog # ASR1397**Specification**

Anti-Horse IgG F(c) Secondary Antibody - Product Information

Description	Anti-HORSE IgG F(c) (GOAT) Antibody
Host	Goat
Conjugate	Unconjugated
Target Species	Horse
Clonality	Polyclonal
Application	WB, E, IC
Application Note	ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000
Physical State	Liquid (sterile filtered)
Host Isotype	IgG
Target Isotype	IgG F(c)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Horse IgG F(c) fragment
Species of Origin	goat
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide

Anti-Horse IgG F(c) Secondary Antibody - Additional Information**Shipping Condition**

Wet Ice

Purity

Anti-Horse IgG F(c) was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Horse IgG, Horse IgG F(c) and Horse Serum. No reaction was observed against Horse IgG F(ab')₂.

Storage Condition

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Horse IgG F(c) Secondary Antibody - Protein Information

Anti-Horse IgG F(c) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Horse IgG F(c) Secondary Antibody - Images

Anti-Horse IgG F(c) Secondary Antibody - Background

Horse IgG F(c) antibody recognizes the F(c) portion of horse IgG. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Horse IgG, Horse IgG F(c) and Horse Serum. No reaction was observed against Horse IgG F(ab')₂. Anti-Horse IgG F(c) antibody is ideal for investigators involved in serum component protein research.