

Bovine IgG F(c)
Catalog # ASR1448**Specification**

Bovine IgG F(c) - Product Information

Description	BOVINE IgG F(c) fragment
Conjugate	Unconjugated
Physical State	Liquid (sterile filtered)
Host Isotype	IgG F(c)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Species of Origin	Bovine
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide

Bovine IgG F(c) - Additional Information**Shipping Condition**

Wet Ice

Purity

This product was prepared from normal serum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and papain digestion followed by chromatographic separation and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Bovine Serum, anti-Bovine IgG and anti-Bovine IgG F(c). No reaction was observed against anti-Bovine IgG F(ab')₂ or anti-Papain.

Storage Condition

Store vial at 4° C prior to restoration. Restore with 1.0 mL of deionized water (or equivalent). For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Bovine IgG F(c) - Protein Information**Bovine IgG F(c) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Bovine IgG F(c) - Images