

Anti-Goat IgG (H&L) Secondary Antibody

Mouse Polyclonal, Unconjugated Catalog # ASR1561

Specification

Anti-Goat IgG (H&L) Secondary Antibody - Product Information

Description Host Conjugate Target Species Reactivity Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen Stabilizer Preservative Anti-GOAT IgG (H&L) (MOUSE) Antibody Mouse Unconjugated Goat Goat Polyclonal WB, E, IC ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000 Liquid (sterile filtered) laG IgG (H&L) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Goat IgG whole molecule None 0.01% (w/v) Sodium Azide

Anti-Goat IgG (H&L) Secondary Antibody - Additional Information

Shipping Condition Wet Ice

Purity

This product was prepared from monospecific polyclonal ascites by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Mouse Serum, Rabbit IgG and Rabbit Serum. No reaction was observed against Goat, Human or Mouse Serum Proteins.

Storage Condition

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Goat IgG (H&L) Secondary Antibody - Protein Information



Anti-Goat IgG (H&L) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Goat IgG (H&L) Secondary Antibody - Images