

# Anti-Horse IgG (H&L) (Rhodamine Conjugated) Secondary Antibody

**Goat Polyclonal, Rhodamine (TRITC)** Catalog # ASR1581

## **Specification**

**Physical State** 

Target Isotype

Host Isotype

Immunogen

**Reconstitution Volume** 

Reconstitution Buffer

Buffer

## Anti-Horse IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Product Information

Anti-HORSE IgG (H&L) (GOAT) Antibody Description

**Rhodamine Conjugated** 

Host Goat **Rhodamine (TRITC)** Conjugate

FP Value 2.4 moles Rhodamine (TRITC) per mole of

> laG Horse

**Target Species** Clonality **Polyclonal Application** IF. FC

**Application Note** FLISA 1:10,000-1:50,000; IF Microscopy

1:1,000-1:5,000;FlowCytometry

1:500-1:2.500 Lyophilized

IgG

IgG (H&L)

0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2 Horse IgG whole molecule

1.0 mL

Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

## Anti-Horse IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Additional **Information**

## **Shipping Condition**

**Ambient** 

### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Horse IgG and Horse Serum.

#### **Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

### **Precautions Note**



Tel: 858.875.1900 Fax: 858.875.1999

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Horse IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Protein Information

### Anti-Horse IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Horse IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Images

Anti-Horse IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.