

Anti-Mouse IgG F(c) Secondary Antibody

Rabbit Polyclonal, Unconjugated Catalog # ASR1619

Specification

Anti-Mouse IgG F(c) Secondary Antibody - Product Information

Anti-MOUSE IgG F(c) (RABBIT) Antibody Description Host Rabbit Conjugate Unconjugated **Target Species** Mouse Reactivity Mouse Clonality Polyclonal Application WB, E, IC **Application Note** ELISA 1:40,000-1:180,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000 **Physical State** Liquid (sterile filtered) Host Isotype laG Target Isotype laG F(c) Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Immunogen Anti-Mouse IgG F(c) was produced by repeated immunization with mouse IgG F(c) fragment in rabbit. Stabilizer None Preservative 0.01% (w/v) Sodium Azide

Anti-Mouse IgG F(c) Secondary Antibody - Additional Information

Shipping Condition Wet Ice

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against Mouse IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Mouse IgG F(c) Secondary Antibody - Protein Information



Anti-Mouse IgG F(c) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Mouse IgG F(c) Secondary Antibody - Images

Anti-Mouse IgG F(c) Secondary Antibody - Background

Anti-Mouse IgG F(c) antibody generated in rabbit detects specifically mouse IgG. This secondary antibody anti-Mouse is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays.