

**F(ab')<sub>2</sub> Anti-Golden Syrian Hamster IgG (H&L) Secondary Antibody**  
**Rabbit Polyclonal, Unconjugated**  
**Catalog # ASR1690****Specification****F(ab')<sub>2</sub> Anti-Golden Syrian Hamster IgG (H&L) Secondary Antibody - Product Information**

Description	<b>F(ab')<sub>2</sub> Anti-GOLDEN SYRIAN HAMSTER IgG (H&amp;L) (RABBIT) Antibody</b>
Host	<b>Rabbit</b>
Conjugate	<b>Unconjugated</b>
Target Species	<b>Golden Syrian Hamster</b>
Clonality	<b>Polyclonal</b>
Application	<b>WB, IHC, E</b>
Application Note	<b>ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunohistochemistry 1:1,000-1:5,000</b>
Physical State	<b>Liquid (sterile filtered)</b>
Host Isotype	<b>IgG F(ab')<sub>2</sub></b>
Target Isotype	<b>IgG (H&amp;L)</b>
Buffer	<b>0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>
Immunogen	<b>Golden Syrian Hamster IgG whole molecule</b>
Stabilizer	<b>None</b>
Preservative	<b>0.01% (w/v) Sodium Azide</b>

**F(ab')<sub>2</sub> Anti-Golden Syrian Hamster IgG (H&L) Secondary Antibody - Additional Information****Shipping Condition**

Wet Ice

**Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Golden Syrian Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Golden Syrian Hamster IgG, Golden Syrian Hamster IgG and Golden Syrian Hamster Serum. No reaction was observed against anti-Pepsin, anti-Rabbit IgG F(c). Limited reactivity will occur with Armenian Hamster IgG.

**Storage Condition**

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

## **F(ab')<sub>2</sub> Anti-Golden Syrian Hamster IgG (H&L) Secondary Antibody - Protein Information**

### **F(ab')<sub>2</sub> Anti-Golden Syrian Hamster IgG (H&L) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **F(ab')<sub>2</sub> Anti-Golden Syrian Hamster IgG (H&L) Secondary Antibody - Images**

### **F(ab')<sub>2</sub> Anti-Golden Syrian Hamster IgG (H&L) Secondary Antibody - Background**

F(ab')<sub>2</sub> Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)<sub>2</sub> fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)<sub>2</sub> fragments penetrate into tissue samples and show better antigen recognition and signal generation in IHC. F(ab)<sub>2</sub> fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')<sub>2</sub> Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.