

## Anti-Cat IgG F(c) Secondary Antibody

Goat Polyclonal, Unconjugated Catalog # ASR1754

### **Specification**

## Anti-Cat IgG F(c) Secondary Antibody - Product Information

Description Anti-CAT IgG F(c) (GOAT) Antibody

Host Goat

Conjugate Unconjugated

Target Species Cat

Clonality Polyclonal Application WB, IHC, E

Application Note ELISA 1:20,000-1:100,000; Western Blot

1:2,000-1:10,000;Immunohistochemistry

1:1,000-1:5,000

Physical State Liquid (sterile filtered)

Host Isotype IgG
Target Isotype IgG F(c)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen Cat IgG F(c) fragment

Stabilizer None

Preservative 0.01% (w/v) Sodium Azide

### Anti-Cat IgG F(c) Secondary Antibody - Additional Information

### **Shipping Condition**

Wet Ice

#### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Cat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Cat IgG, Cat IgG F(c) and Cat Serum. No reaction was observed against Cat IgG F(ab')2.

### **Storage Condition**


Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

### Anti-Cat IgG F(c) Secondary Antibody - Protein Information

# Anti-Cat IgG F(c) Secondary Antibody - Protocols





Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Cat IgG F(c) Secondary Antibody - Images