

Anti-Human IgA (alpha chain) (Biotin Conjugated) Secondary Antibody

Goat Polyclonal, Biotin Catalog # ASR1779

Specification

Anti-Human IgA (alpha chain) (Biotin Conjugated) Secondary Antibody - Product Information

Description Anti-HUMAN IgA (alpha chain) (GOAT)

Antibody Biotin Conjugated

Host Goat Conjugate Biotin

FP Value 10-20 moles Biotin per mole of IgG

Target Species
Reactivity
Human
Clonality
Application
Human
Polyclonal
WB, E, IC

Application Note ELISA 1:60,000-1:600,000; Western Blot

1:2,000-1:10,000;Immunochemistry

1:1,000-1:5,000 Lyophilized

Physical State
Host Isotype
Target Isotype
IgA

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen Anti-Human IgA alpha heavy chain

antibody was produced by repeated

immunization with Human IgA alpha heavy

chain in goat.

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Anti-Human IgA (alpha chain) (Biotin Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

Anti-HUMAN IgA (alpha chain) (GOAT) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgA coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Human IgA and Human Serum. No reaction was observed against other Human heavy or light chain proteins.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C



Tel: 858.875.1900 Fax: 858.875.1999

or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Human IgA (alpha chain) (Biotin Conjugated) Secondary Antibody - Protein Information

Anti-Human IgA (alpha chain) (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cvtometv
- Cell Culture

Anti-Human IgA (alpha chain) (Biotin Conjugated) Secondary Antibody - Images Anti-Human IgA (alpha chain) (Biotin Conjugated) Secondary Antibody - Background

Anti-Human IgA alpha heavy chain antibody generated in goat detects specifically Human IgA alpha heavy chain. This secondary antibody anti-Human is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays.