

Anti-Human IgG F(c) (Rhodamine Conjugated) Secondary Antibody

Rabbit Polyclonal, Rhodamine (TRITC) Catalog # ASR1787

Specification

Physical State Host Isotype

Target Isotype

Immunogen

Reconstitution Volume

Reconstitution Buffer

Buffer

Anti-Human IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Product Information

Anti-HUMAN IgG F(c) (RABBIT) Antibody Description

Rhodamine Conjugated

Host Rabbit **Rhodamine (TRITC)** Conjugate

FP Value 1.5 moles Rhodamine (TRITC) per mole of

> Human **Polyclonal**

Target Species Clonality **Application** IF. FC

Application Note FLISA 1:10,000-1:50,000; IF Microscopy

1:1,000-1:5,000;FlowCytometry

1:500-1:2.500 Lyophilized IgG

IgG F(c)

0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2 **Human IgG F(c) fragment**

1.0 mL

Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Anti-Human IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Human IgG, Human IgG F(c) and Human Serum. No reaction was observed against Human IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note



Tel: 858.875.1900 Fax: 858.875.1999

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Human IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Protein Information

Anti-Human IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Human IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Images

Anti-Human IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.