

Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody

Goat Polyclonal, Rhodamine (TRITC) Catalog # ASR1794

Specification

Physical State

Immunogen

Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Product Information

Anti-MOUSE IgG F(c) (GOAT) Antibody Description

Rhodamine Conjugated

Host Goat **Rhodamine (TRITC)** Conjugate

FP Value 3.9 moles Rhodamine (TRITC) per mole of

> Mouse **Polyclonal**

Target Species Clonality **Application** IF. FC

Application Note FLISA 1:10,000-1:50,000; IF Microscopy

1:1,000-1:5,000;FlowCytometry

1:500-1:2.500 Lyophilized

Host Isotype IgG Target Isotype IgG F(c)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2 Mouse IgG F(c) fragment

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against Mouse IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note



Tel: 858.875.1900 Fax: 858.875.1999

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Protein Information

Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Images

Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.