

## Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody

Rabbit Polyclonal, Rhodamine (TRITC) Catalog # ASR1804

#### Specification

### Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Product Information

Description

Host Conjugate FP Value

Target Species Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen Reconstitution Volume Reconstitution Buffer

Stabilizer

Preservative

Anti-MOUSE IgG F(c) (RABBIT) Antibody **Rhodamine Conjugated** Rabbit **Rhodamine (TRITC)** 2.3 moles Rhodamine (TRITC) per mole of laG Mouse Polyclonal IF. FC FLISA 1:10,000-1:50,000;IF Microscopy 1:1,000-1:5,000;FlowCytometry 1:500-1:2.500 Lyophilized lgG lgG F(c) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Mouse IgG F(c) fragment 1.0 mL Restore with deionized water (or equivalent) 10 mg/mL Bovine Serum Albumin (BSA) -Immunoglobulin and Protease free 0.01% (w/v) Sodium Azide

# Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Additional Information

Shipping Condition Ambient

#### Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against Mouse IgG F(ab')2.

#### Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.



#### Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

#### Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Protein Information

#### Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

#### Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Images

#### Anti-Mouse IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.